

A perturbative probabilistic approach to quantum many-body systems

Andrea Di Stefano^{1,2}, Massimo Ostilli^{1,3} and Carlo Presilla^{1,4}

¹ Dipartimento di Fisica, Università di Roma “La Sapienza”, Piazzale A. Moro 2, Roma 00185, Italy

² Fakultät für Mathematik, Universität Bielefeld, Postfach 110 131, 33501 Bielefeld, Germany

³ Cooperative Association for Internet Data Analysis (CAIDA), University of California, San Diego (UCSD), 9500 Gilman Drive, La Jolla, California 92093, USA

⁴ Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Roma 1, Roma 00185, Italy

Abstract. In the probabilistic approach to quantum many-body systems, the ground-state energy is the solution of a nonlinear scalar equation written either as a cumulant expansion or as an expectation with respect to a probability distribution of the potential and hopping (amplitude and phase) values recorded during an infinitely lengthy evolution. We introduce a perturbative expansion of this probability distribution which conserves, at any order, a multinomial-like structure, typical of uncorrelated systems, but includes, order by order, the statistical correlations provided by the cumulant expansion. The proposed perturbative scheme is successfully tested in the case of spin $\frac{1}{2}$ hard-core boson Hubbard models.

PACS numbers: 02.50.-r, 05.40.-a, 71.10.Fd

1. Introduction

A multitude of evolution problems, including quantum many-body systems, can be cast in the form of a linear flow, namely a system of linear differential equations with respect to a parameter, the time, which can be real or imaginary. The solution of a linear flow, with a real or an imaginary time, admits an exact probabilistic representation, namely a Feynman-Kac-like formula, in terms of a proper collection of independent Poisson processes [1, 2, 3, 4, 5]. For a lattice system, the Poisson processes are associated with the links of the lattice and the probabilistic representation leads to an optimal algorithm [3, 4, 5] which coincides with the Green function quantum Monte Carlo method in the limit when the latter becomes exact [6]. The algorithm can be rigorously generalized [7] to include fluctuation control techniques, like reconfigurations and importance sampling, and allows the exact simulation of time-dependent correlation functions for system not affected by the so called sign problem [8].

In the limit of an infinitely-long imaginary time, the above exact probabilistic representation has been developed to yield semi-analytical results. In fact, for an arbitrary many-body system we are able to relate the energy of its ground state to the unique solution of a nonlinear scalar equation [9, 10, 11, 12]. This equation can be written in terms of a series involving the cumulants of integers, the multiplicities N_V , N_T and N_λ , which count how many times the potential, hopping and phase variables take the values V , T and λ during an infinitely long evolution of the system. The potential variables are, in some chosen base, the diagonal matrix elements of the Hamiltonian of the system, whereas the hopping and phase variables are related to the amplitude and phase of the off-diagonal matrix elements. Alternatively, the equation for the ground-state energy can be written in terms of an expectation involving the probability distribution of the multiplicities N_V , N_T and N_λ .

The two ways of writing the equation for E_0 , cumulant expansion or probabilistic expectation, correspond to two different approaches to evaluate the ground state of a many-body system. In the former case, we can imagine to measure the exact cumulants of the system up to some finite (small) order, insert them in a corresponding truncated equation and solve it obtaining an approximation to E_0 [11]. In the latter case, we have the possibility to make some guess on the probability distribution of the multiplicities N_V , N_T and N_λ bypassing the microscopic connection between configurations of the system and values of the variables V , T and λ . The simplest guess is to neglect any correlation among the multiplicities and assume that they are multinomially distributed. There is a class of systems for which, in the thermodynamic limit, a multinomial distribution exactly applies. The class includes the uniformly fully connected models, namely a collection of states all connected with equal hopping coefficients and in the presence of a potential operator with arbitrary levels and degeneracies, and the random potential systems, in which the hopping operator is generic and arbitrary potential levels are assigned randomly to the states with arbitrary probabilities. For this class of models we have found a zero-temperature universal thermodynamic limit displaying a quantum

phase transition [12].

Both the two approaches described above have limitations. The more severe drawback of the cumulant expansion is that truncating the series corresponds to introducing a rather artificial equivalent system with zero cumulants of large order. Since the determination of E_0 is a problem which involves large fluctuations [9], we expect, and find indeed, some nonphysical behavior of the solutions corresponding to large interaction energies. On the other hand, we expect that the multinomial probability distribution used in the second approach may give quantitatively inaccurate results for most of the systems, i.e. when the correlations among the potential, hopping and phase multiplicities cannot be neglected.

In the present paper we introduce a perturbative scheme merging the merits of the cumulant expansion with those of the expectation taken from an uncorrelated multinomial distribution. The idea is to develop a perturbative expansion of the probability distribution of the multiplicities N_V , N_T and N_λ which conserves, at any order, a multinomial-like structure. Order by order, we add correlations among the multiplicities in such a way to modify the cumulants of the multinomial-like distribution and make them to coincide with those measured in the system up to the order considered. As a result, we gain better and better approximations to the real probability distribution. At the first order, we have a probability distribution which contain infinitely many cumulants and the first one is exact; at the second order, we have a probability distribution with infinitely many cumulants and the first two are exact; and so on. In this way, we expect to obtain, even at very small perturbative orders, accurate results for the ground-state energy of both weakly- and strongly-interacting many-body systems. We have checked our perturbative scheme in the case of spin $\frac{1}{2}$ hard core boson Hubbard models in two dimensional square lattices. It is remarkable that, already at the second perturbative order, we find a ground-state energy which compares rather well with the exact value of E_0 .

The main advantage of our method lies in its semi-analytical character. Once all the cumulants up to some order k are measured, via a *una tantum* simulation, the perturbative probabilistic distribution built from these input data provides, within an approximation which improves with k , the ground state energy E_0 as a function of the Hamiltonian parameters (coupling and hopping amplitudes). In contrast, in a standard Monte Carlo method any different choice of the Hamiltonian parameters requires a distinct simulation. Furthermore, the cumulants are easily measured also in the case of fermions (bosons in the presence of magnetic fields). The sign (phase) problem that occurs in this case remains confined in the expression of the probability distribution and can, in principle, be addressed analytically.

The paper is organized as follows. In section 2 we review the probabilistic approach to quantum many-body systems. The equation for the ground-state energy is written as an expansion over the cumulants of the potential, hopping and phase multiplicities in subsection 2.1 and as an expectation with respect to the probability distribution of the same variables in subsection 2.2. In the latter subsection the case of an

uncorrelated multinomial distribution is described in detail. In section 3 we introduce the probabilistic perturbative scheme bringing together the merits of the multinomial probability distribution with the statistical details provided by the cumulant expansion. The parameters defining the perturbative probability distribution of the potential, hopping and phase multiplicities are explicitly discussed, up to the third order, in subsections from 3.1 to 3.5. This is a technical part which could be skipped in a first reading. Some considerations on the higher perturbative orders are given in subsection 3.6. The equation for the evaluation of E_0 resulting from the above perturbative scheme is discussed in section 4. Section 5 deals with some numerical results. Concluding remarks are drawn in section 6. Two appendices close the paper. In Appendix A we review the methods for solving the nonsymmetric algebraic Riccati equation which appears in determining the perturbative parameters at the second order. Appendix B summarizes the evaluation of the perturbative parameters at the fourth order.

2. Probabilistic approach to quantum many-body systems

In this section we review the probabilistic approach to quantum many-body systems developed in [9, 10, 11, 12]. Consider a system of particles represented by a Hamiltonian operator \hat{H} acting on a M dimensional space of states labeled by configuration indexes \mathbf{n} . As an example, think about spinless particles undergoing a simple exclusion dynamics in a lattice with configurations given by the site occupations $\mathbf{n} = (0, 1, 1, 1, 0, 0, 1, \dots)$. In the chosen \mathbf{n} -representation, we separate, as usual, $\hat{H} = \hat{K} + \hat{V}$, where $\hat{V} = \text{diag}(\hat{H})$ is called the potential operator and has non vanishing matrix elements

$$V_{\mathbf{n},\mathbf{n}} = V_{\mathbf{n}}. \quad (1)$$

The hopping operator \hat{K} is defined by the matrix elements

$$K_{\mathbf{n},\mathbf{n}'} = -\lambda_{\mathbf{n},\mathbf{n}'} \eta_{\mathbf{n},\mathbf{n}'}, \quad \eta_{\mathbf{n},\mathbf{n}'} > 0, \quad |\lambda_{\mathbf{n},\mathbf{n}'}| = 0, 1, \quad (2)$$

conventionally written in terms of links $\lambda_{\mathbf{n},\mathbf{n}'}$ and positive strengths $\eta_{\mathbf{n},\mathbf{n}'}$. The matrix with elements $|\lambda_{\mathbf{n},\mathbf{n}'}|$ forms the so called adjacency matrix and establishes whether two configurations \mathbf{n} and \mathbf{n}' are first neighbours with respect to \hat{K} or not. The phase change (sign in the case of fermions) registered by connecting two first neighbour configurations \mathbf{n} and \mathbf{n}' is registered by $\text{Arg } \lambda_{\mathbf{n},\mathbf{n}'}$. Note that $\sum_{\mathbf{n}'} |\lambda_{\mathbf{n},\mathbf{n}'}|$ is the number of configurations which may be connected to the configuration \mathbf{n} .

The evolution of the system from an initial state ψ_0 is earned by solving the Schrödinger equation. After a time t the system is found in the state $\psi(t)$ which in the chosen \mathbf{n} -representation has components (we put $\hbar = 1$)

$$\langle \mathbf{n}, \psi(t) \rangle = \sum_{\mathbf{n}_0} \langle \mathbf{n}, e^{-i\hat{H}t} \mathbf{n}_0 \rangle \langle \mathbf{n}_0, \psi_0 \rangle. \quad (3)$$

This evolution, as any linear flow, admits an exact probabilistic representation [1, 2, 3, 4, 5]. In a one-to-one correspondence with the links, we introduce M^2 independent Poisson processes $\{N_{\mathbf{n},\mathbf{n}'}^t\}$ with rates $\{\rho_{\mathbf{n},\mathbf{n}'}\}$. We recall that for these



Figure 1. Random walk with N_t jumps in the time interval $[0, t)$: scheme of the visited configurations and of the corresponding jump times.

processes the probability to jump k times in the time interval $[t, t + s)$ is [13]

$$\text{prob}(N_{\mathbf{n}, \mathbf{n}'}^{t+s} - N_{\mathbf{n}, \mathbf{n}'}^t = k) = \frac{(\rho_{\mathbf{n}, \mathbf{n}'} s)^k}{k!} e^{-\rho_{\mathbf{n}, \mathbf{n}'} s}, \quad k = 0, 1, 2, \dots \quad (4)$$

We establish that each time a Poisson process $N_{\mathbf{n}, \mathbf{n}'}^t$ jumps, the configuration of the system changes from \mathbf{n} to \mathbf{n}' if $|\lambda_{\mathbf{n}, \mathbf{n}'}| = 1$, otherwise remains \mathbf{n} . Arranging the jumps according to the times, $s_1 < s_2 < \dots < s_{N_t} < t < s_{N_t+1}$, at which they take place in the interval $[0, t)$ we define a random walk in the configuration space of the system $\mathbf{n}_0 \rightarrow \mathbf{n}_1 \rightarrow \mathbf{n}_2 \rightarrow \dots \rightarrow \mathbf{n}_{N_t}$ generated by the above rule from a chosen initial configuration \mathbf{n}_0 . It is simple to prove that the fundamental matrix elements of the evolution operator $\langle \mathbf{n}, e^{-i\hat{H}t} \mathbf{n}_0 \rangle$ can be written as an expectation over the above M^2 independent Poisson processes [14]

$$\langle \mathbf{n}, e^{-i\hat{H}t} \mathbf{n}_0 \rangle = \mathbb{E}(\delta_{\mathbf{n}, \mathbf{n}_{N_t}} \mathcal{M}_{\mathbf{n}_0}^t), \quad (5)$$

$$\mathcal{M}_{\mathbf{n}_0}^t = e^{\sum_{\mathbf{n}, \mathbf{n}'} \rho_{\mathbf{n}, \mathbf{n}'} t} \left(\prod_{k=1}^{N_t} i\lambda_k \eta_k \rho_k^{-1} e^{-iV_{k-1}(s_k - s_{k-1})} \right) e^{-iV_{N_t}(t - s_{N_t})}, \quad (6)$$

where we put $s_0 = 0$ and introduced the shorthand

$$V_k = V_{\mathbf{n}_k}, \quad k = 0, 1, \dots, N_t, \quad (7)$$

$$\lambda_k = \lambda_{\mathbf{n}_{k-1}, \mathbf{n}_k}, \quad k = 1, \dots, N_t, \quad (8)$$

$$\eta_k = \eta_{\mathbf{n}_{k-1}, \mathbf{n}_k}, \quad \rho_k = \rho_{\mathbf{n}_{k-1}, \mathbf{n}_k}, \quad k = 1, \dots, N_t. \quad (9)$$

The rates $\{\rho_{\mathbf{n}, \mathbf{n}'}\}$ of the Poisson processes are completely arbitrary, in fact it is easy to check that $\mathbb{E}(d\mathcal{M}_{\mathbf{n}_0}^t/d\rho_{\mathbf{n}, \mathbf{n}'}) = 0$. Here, for simplicity, we take $\rho_{\mathbf{n}, \mathbf{n}'} = \rho$ uniform, whereas other choices, e.g. $\rho_{\mathbf{n}, \mathbf{n}'} = \eta_{\mathbf{n}, \mathbf{n}'}$, allow to define optimal Monte Carlo numerical algorithms [3, 7]. The above probabilistic representation holds also for non-autonomous systems with a time dependent potential $V_{\mathbf{n}}(t)$. In this case the time-ordered quantum evolution operator $\mathbb{T} \exp(-i \int_0^t \hat{H}(u) du)$ has matrix elements given by (5-6) with $V_k \rightarrow \int_{s_k}^{s_{k+1}} V_k(u) du$. The representation holds also at imaginary times $t \rightarrow -it$ with the substitutions $\lambda_k \rightarrow -i\lambda_k$ and $V_k \rightarrow -iV_k$.

A convenient way to study the properties of the ground state of a particle system is to consider its evolution for a long imaginary time. Starting from an arbitrary configuration \mathbf{n}_0 the system finally relaxes into the ground state, assumed not to be orthogonal to \mathbf{n}_0 . In this way we can evaluate any ground-state correlation function via time asymptotic probabilistic expressions. For instance, the ground state energy E_0 is given by

$$E_0 = \lim_{t \rightarrow \infty} -\partial_t \log \sum_{\mathbf{n}} \langle \mathbf{n}, e^{-\hat{H}t} \mathbf{n}_0 \rangle = \lim_{t \rightarrow \infty} -\partial_t \log \mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t), \quad (10)$$

where $\mathcal{M}_{\mathbf{n}_0}^t$ is the imaginary time variant of (6). It is of fundamental importance that for the expectation $\mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t)$ we can find, at large times, an analytical expression. In the following we outline how this result is obtained.

First, we decompose the expectation in a series of canonical expectations over random walks with a fixed number of jumps

$$\mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t) = \sum_{N=0}^{\infty} \mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t, N_t = N) \quad (11)$$

and evaluate each term of this series by integrating over all possible jump times. The probability to have N Poisson processes jumping in the interval $[0, t)$ with the k -th process jumping in the interval $[s_k, s_k + ds_k)$ amounts to

$$\prod_{k=1}^N e^{-\sum_{\mathbf{n}, \mathbf{n}'} \rho_{\mathbf{n}, \mathbf{n}'}(s_k - s_{k-1})} \rho_k ds_k. \quad (12)$$

With a simple calculation we then obtain

$$\mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t, N_t = N) = \sum_{r \in \Omega_N} \mathcal{W}_N^{(r)}(t) \prod_{k=1}^N \lambda_k^{(r)} \eta_k^{(r)}, \quad (13)$$

where $\Omega_N = \Omega_N(\mathbf{n}_0)$ is the set of all possible random walks with N jumps branching from \mathbf{n}_0 . The contribution of the r -th random walk $\mathbf{n}_0 \rightarrow \mathbf{n}_1^{(r)} \rightarrow \dots \rightarrow \mathbf{n}_N^{(r)}$ includes the factor $\mathcal{W}_N^{(r)}(t) = \mathcal{L}^{-1}[\tilde{\mathcal{W}}_N^{(r)}(z)](t)$ which is the inverse Laplace transform of

$$\tilde{\mathcal{W}}_N^{(r)}(z) = \prod_{k=1}^N \frac{1}{z + V_k^{(r)}}. \quad (14)$$

From equation (13) it is evident that only the random walks with $|\lambda_k^{(r)}| = 1$, i.e. $\mathbf{n}_{k-1}^{(r)} \neq \mathbf{n}_k^{(r)}$, for $k = 1, \dots, N$, contribute to the sum. Thus we can rewrite the sum over Ω_N as a probabilistic expectation over these effective random walks. According to the choice $\rho_{\mathbf{n}, \mathbf{n}'} = \rho$ uniform, the effective random walks correspond to a Markov chain with transition matrix $P_{\mathbf{n}, \mathbf{n}'} = |\lambda_{\mathbf{n}, \mathbf{n}'}| / \sum_{\mathbf{n}''} |\lambda_{\mathbf{n}, \mathbf{n}''}|$ [15]. The probability that we must associate to the r -th element of the set Ω_N is, therefore,

$$p_N^{(r)} = \prod_{k=1}^N \frac{|\lambda_{\mathbf{n}_{k-1}^{(r)}, \mathbf{n}_k^{(r)}}|}{\sum_{\mathbf{n}} |\lambda_{\mathbf{n}_{k-1}^{(r)}, \mathbf{n}}|}. \quad (15)$$

Note that $p_N^{(r)} = 0$ for non effective random walks and $\sum_{r \in \Omega_N} p_N^{(r)} = 1$. Multiplying and dividing by $p_N^{(r)}$, we rewrite the canonical expectations as

$$\mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t, N_t = N) = \sum_{r \in \Omega_N} p_N^{(r)} \mathcal{L}^{-1} \left[\prod_{k=1}^N \frac{1}{z + V_k^{(r)}} \right] (t) \prod_{k=1}^N T_k^{(r)} \prod_{k=1}^N \lambda_k^{(r)}, \quad (16)$$

where

$$T_k = \eta_{\mathbf{n}_{k-1}, \mathbf{n}_k} \sum_{\mathbf{n}} |\lambda_{\mathbf{n}_{k-1}, \mathbf{n}}|, \quad k = 1, \dots, N. \quad (17)$$

Equation (16) shows that $\mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t, N_t = N)$ is the average of a quantity which does not rely on the detailed sequence of the configurations visited during the time t . It

depends just on the multiplicities, or numbers of occurrences, of the potential, hopping and phase variables, V , T and λ , defined by (7), (17) and (8), respectively. For a random walk with N jumps, these multiplicities are explicitly defined as

$$N_V = \sum_{k=0}^N \delta_{V,V_k}, \quad V \in \mathcal{V}, \quad (18)$$

$$N_T = \sum_{k=1}^N \delta_{T,T_k}, \quad T \in \mathcal{T}, \quad (19)$$

$$N_\lambda = \sum_{k=1}^N \delta_{\lambda,\lambda_k}, \quad \lambda \in \mathcal{L}, \quad (20)$$

where \mathcal{V} , \mathcal{T} and \mathcal{L} are the sets of all possible values assumed by (7), (17) and (8) during a random walk with infinitely many jumps. Note that $0 \notin \mathcal{L}$ as jumps between configurations \mathbf{n} and \mathbf{n}' with $\lambda_{\mathbf{n},\mathbf{n}'} = 0$ have zero probability to be realized. Since any configuration can be obtained from any other one by a finite number of jumps, i.e. the Markov chain we are considering is irreducible, the elements in the sets \mathcal{V} , \mathcal{T} and \mathcal{L} do not depend on the initial configuration \mathbf{n}_0 . Let us indicate the multiplicities N_V , N_T and N_λ collectively by a vector with as many components as the elements in the set $\mathcal{H} = \mathcal{V} \cup \mathcal{T} \cup \mathcal{L}$,

$$\boldsymbol{\mu}^T = (\dots N_V \dots; \dots N_T \dots; \dots N_\lambda \dots). \quad (21)$$

If we split the set Ω_N into subsets of random walks with equal values of $\boldsymbol{\mu}$, we conclude that

$$\begin{aligned} \mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t, N_t = N) = \\ \sum_{\boldsymbol{\mu}} \mathcal{P}_N(\boldsymbol{\mu}) \mathcal{L}^{-1} \left[\prod_{V \in \mathcal{V}} (z + V)^{-N_V} \right] (t) \prod_{T \in \mathcal{T}} T^{N_T} \prod_{\lambda \in \mathcal{L}} \lambda^{N_\lambda}, \end{aligned} \quad (22)$$

where

$$\mathcal{P}_N(\boldsymbol{\mu}) = \sum_{r \in \Omega_N} \left(\prod_{k=1}^N \frac{|\lambda_{\mathbf{n}_{k-1}^{(r)}, \mathbf{n}_k^{(r)}}|}{\sum_{\mathbf{n}} |\lambda_{\mathbf{n}_{k-1}^{(r)}, \mathbf{n}}|} \right) \delta_{\boldsymbol{\mu}^{(r)}, \boldsymbol{\mu}} \quad (23)$$

is the probability to have random walks with multiplicities $\boldsymbol{\mu}$ after N jumps from the configuration \mathbf{n}_0 . Note that $\mathcal{P}_N(\boldsymbol{\mu}) = 0$ unless $\boldsymbol{\mu}$ satisfies the following three constraints

$$\sum_{V \in \mathcal{V}} N_V = N + 1, \quad \sum_{T \in \mathcal{T}} N_T = N, \quad \sum_{\lambda \in \mathcal{L}} N_\lambda = N. \quad (24)$$

As a second step, we observe that when the imaginary time t becomes large the full expectation (11) takes exponentially leading contributions from terms with $N \sim t$. Thus, we can replace all previous results with their $N \rightarrow \infty$ asymptotic expressions. Due to the ergodicity of the underlying Markov chain, the probability (23) loses memory of the initial configuration \mathbf{n}_0 . We can evaluate the inverse Laplace transform which appears in (22) by a saddle point technique in the complex plane. In the same equation we can also substitute the sum over the multiplicities by an integral over $\boldsymbol{\mu}$ and avoid

to distinguish the normalizations N and $N + 1$ in (24). As a final result we have the following asymptotic logarithm equality

$$\mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t, N_t = N) \simeq \int d(N\boldsymbol{\nu}) \mathcal{P}_N(N\boldsymbol{\nu}) \frac{e^{x_0 t + N(\boldsymbol{\nu}, \mathbf{u})}}{\sqrt{2\pi N(\boldsymbol{\nu}, \mathbf{w})}}, \quad (25)$$

where we have introduced the frequencies $\boldsymbol{\nu} = \boldsymbol{\mu}/N$, which have a finite limit for $N \rightarrow \infty$, the vectors

$$\mathbf{u}^T = (\dots - \log(x_0 + V) \dots; \dots \log T \dots; \dots \log \lambda \dots), \quad (26)$$

$$\mathbf{w}^T = (\dots (x_0 + V)^{-2} \dots; \dots 0 \dots; \dots 0 \dots), \quad (27)$$

and the scalar product $(\mathbf{a}, \mathbf{b}) = \sum_{\alpha \in \mathcal{H}} a_\alpha b_\alpha$. The quantity x_0 , which is the real saddle-point in the complex contour used to evaluate the Laplace antitransform, is the unique solution of the scalar equation

$$\sum_{V \in \mathcal{V}} \frac{\nu_V}{x_0 + V} = \frac{t}{N}, \quad x_0 > -V_{\min}. \quad (28)$$

Note that this equation has a regular scaling behavior for $t, N \rightarrow \infty$ with $N \sim t$.

The integral over the frequencies $\boldsymbol{\nu}$ in equation (25) is easily performed by a saddle-point method (actually Laplace's method) if the asymptotic probability density $\mathcal{P}_N(N\boldsymbol{\nu})$ is known. We cannot hope to evaluate this probability from the microscopic definition (23) except for very particular models. For general systems, two different strategies have been considered, see [11] and [12]. We can relate $\mathcal{P}_N(N\boldsymbol{\nu})$ to proper statistical moments of the system under consideration, measure or calculate some of these moments and, lastly, obtain partial information about E_0 . Or we can postulate some expressions for $\mathcal{P}_N(N\boldsymbol{\nu})$ and see which kind of systems are described, exactly or approximately, by this guess.

2.1. Cumulant expansion

If we rewrite the probability density $\mathcal{P}_N(N\boldsymbol{\nu})$ in terms of its Fourier transform $\tilde{\mathcal{P}}_N(\mathbf{q})$

$$\mathcal{P}_N(N\boldsymbol{\nu}) = (2\pi)^{-|\mathcal{H}|} \int d\mathbf{q} \tilde{\mathcal{P}}_N(\mathbf{q}) e^{-i(\mathbf{q}, N\boldsymbol{\nu})}, \quad (29)$$

we can associate $\log \tilde{\mathcal{P}}_N(\mathbf{q})$ with the cumulants, or connected correlation functions, of the multiplicities $N\boldsymbol{\nu}$ sampled with respect to the measure $\mathcal{P}_N(N\boldsymbol{\nu})$. Indicating with $\langle N\nu_{\alpha_1} \dots N\nu_{\alpha_k} \rangle_N^{(c)}$ the component $\alpha_1 \dots \alpha_k$ of the cumulant of order k , we have the well known relation [16]

$$\begin{aligned} \log \tilde{\mathcal{P}}_N(\mathbf{q}) &= \sum_{k=1}^{\infty} \frac{1}{k!} \langle (N\boldsymbol{\nu}, i\mathbf{q})^k \rangle_N^{(c)} \\ &= \sum_{k=1}^{\infty} \frac{i^k}{k!} \sum_{\alpha_1 \in \mathcal{H}} \dots \sum_{\alpha_k \in \mathcal{H}} \langle N\nu_{\alpha_1} \dots N\nu_{\alpha_k} \rangle_N^{(c)} q_{\alpha_1} \dots q_{\alpha_k}. \end{aligned} \quad (30)$$

At this point, it is simple to calculate the canonical expectations $\mathbb{E}(\mathcal{M}_{\mathbf{n}_0}^t, N_t = N)$ performing the integrals over the $2|\mathcal{H}|$ variables $\boldsymbol{\nu}$ and \mathbf{q} by a saddle-point

approximation. This approximation is asymptotically exact for $N \rightarrow \infty$. The last step is to evaluate E_0 by resumming the series (11). By virtue of the limit $t \rightarrow \infty$ to be taken at the end, we still have an exact result if we replace the series over N by an integral and estimate this integral at its saddle-point. There is one important point to observe. Independently of their order k , the cumulants $\langle N\nu_{\alpha_1} \dots N\nu_{\alpha_k} \rangle_N^{(c)}$ diverge as N for $N \rightarrow \infty$. Thus we introduce the asymptotic rescaled cumulants

$$\Sigma_{\alpha_1, \dots, \alpha_k}^{(k)} = \lim_{N \rightarrow \infty} \frac{1}{N} \langle N\nu_{\alpha_1} \dots N\nu_{\alpha_k} \rangle_N^{(c)}, \quad (31)$$

$\Sigma^{(k)}$ in a compact notation. Existence and finiteness of these limits are ensured by the finite correlation length N_c which characterizes the correlations functions, connected or not, of the the multiplicities [11]. We conclude that E_0 is the unique solution of the scalar equation

$$\sum_{k=1}^{\infty} \frac{1}{k!} \sum_{\alpha_1 \in \mathcal{H}} \dots \sum_{\alpha_k \in \mathcal{H}} \Sigma_{\alpha_1, \dots, \alpha_k}^{(k)} u_{\alpha_1}(E_0) \dots u_{\alpha_k}(E_0) = 0, \quad E_0 \leq V_{\min}, \quad (32)$$

where

$$\mathbf{u}^T(E_0) = (\dots - \log(-E_0 + V) \dots; \dots \log T \dots; \dots \log \lambda \dots). \quad (33)$$

Equation (32) is exact and the uniqueness of its solution is ensured by the constraint $E_0 \leq V_{\min}$ which stems from the Laplace transform causality condition (28).

To find the exact ground-state energy E_0 from equation (32) we have to know the cumulants $\Sigma^{(k)}$ at any order k . Of course, for a general system this is not conceivable. However, up to some small order k and even for systems of relatively large size, the cumulants can be measured by reliable statistical simulations [11]. With these input data, we can truncate the series in (32) at some order and obtain an approximation to E_0 . Independently of the truncation order, there is an important feature to note. Suppose that we change the Hamiltonian \hat{H} leaving unaltered the adjacency matrix $|\lambda_{\mathbf{n}, \mathbf{n}'}|$ and the number of elements in the sets \mathcal{V} , \mathcal{T} and \mathcal{L} . The asymptotic rescaled cumulants $\Sigma^{(k)}$ are unaffected by this change and the only modifications to equation (32) are encoded analytically by $\mathbf{u}(E_0)$. Therefore, the same input data $\Sigma^{(k)}$, $k = 1, \dots, k_{\max}$, can be used to find the ground-state energy of parametric Hamiltonians as a function of their parameters. The simplest example is to consider $\hat{H}(\gamma) = \hat{K} + \gamma \hat{V}$ and evaluate the function $E_0(\gamma)$.

At the lowest truncation order $k_{\max} = 1$, equation (32) reads

$$\sum_{V \in \mathcal{V}} \Sigma_V^{(1)} \log(-E_0 + V) = \sum_{T \in \mathcal{T}} \Sigma_T^{(1)} \log T + \sum_{\lambda \in \mathcal{L}} \Sigma_{\lambda}^{(1)} \log \lambda, \quad E_0 \leq V_{\min}. \quad (34)$$

In general this equation must be solved numerically. Conversely, for $\hat{V} = 0$ we have $\mathcal{V} = \{0\}$ and $\Sigma_{V=0}^{(1)} = 1$ which allow us to find the analytical solution

$$E_0^{(0)} = - \left(\prod_{T \in \mathcal{T}} T^{\Sigma_T^{(1)}} \right) \left(\prod_{\lambda \in \mathcal{L}} \lambda^{\Sigma_{\lambda}^{(1)}} \right). \quad (35)$$

This represents the lowest order approximation to the ground-state energy of the hopping operator \hat{K} .

The cumulant expansion described above has been applied to study Hubbard models in a two dimensional lattice [11]. By including cumulants up to order 4, the results compare rather well with the exact ground-state energy (determined numerically in other ways) at least in the case of hard core bosons, i.e. $\mathcal{L} = \{1\}$, and for interaction energies not too large with respect to the hopping term. The results, however, are disappointing for large interaction energies, in fact in this limit E_0 diverges. Moreover, in the case of fermions when $\mathcal{L} = \{-1, 1\}$ pointless complex solutions can be found for E_0 as shown, for instance, for $\hat{V} = 0$ and at the lowest order by equation (35). These problems stem from the fact that truncating at some order k_{\max} equation (32) is equivalent to consider an artificial probability density $\mathcal{P}_N(N\boldsymbol{\nu})$ with zero cumulants at any order $k > k_{\max}$.

2.2. Multinomial probability density

In view of the definition (31) of the asymptotic rescaled cumulants, equation (32) can be compactly rewritten as

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^{\infty} \frac{1}{k!} \langle (N\boldsymbol{\nu}, \mathbf{u}(E_0))^k \rangle_N^{(c)} = 0, \quad E_0 \leq V_{\min}. \quad (36)$$

The series sums up to $\log \tilde{\mathcal{P}}_N(-i\mathbf{u}(E_0))$ hence we can state that E_0 is the unique solution of the exact equation

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log \int d(N\boldsymbol{\nu}) \mathcal{P}_N(N\boldsymbol{\nu}) e^{(N\boldsymbol{\nu}, \mathbf{u}(E_0))} = 0, \quad E_0 \leq V_{\min}, \quad (37)$$

with $\mathbf{u}(E_0)$ given by (33). Equation (37) makes crystal clear that the knowledge of E_0 stems from that of $\mathcal{P}_N(N\boldsymbol{\nu})$.

The random walks in the configuration space definitely induce correlations among the multiplicities $\boldsymbol{\mu}$ due to the dependence of the potential, hopping and phase variables on the visited configurations. Could we neglect these correlations, for N sufficiently large the multiplicities of each set \mathcal{V} , \mathcal{T} and \mathcal{L} would be equivalent to multinomial trials processes with success probabilities

$$p_V = \lim_{N \rightarrow \infty} \frac{1}{N+1} \sum_{k=0}^N \delta_{V, V_k} = \Sigma_V^{(1)}, \quad V \in \mathcal{V}, \quad (38)$$

$$p_T = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N \delta_{T, T_k} = \Sigma_T^{(1)}, \quad T \in \mathcal{T}, \quad (39)$$

$$p_\lambda = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N \delta_{\lambda, \lambda_k} = \Sigma_\lambda^{(1)}, \quad \lambda \in \mathcal{L}. \quad (40)$$

Note that we made use of the ergodic properties of the underlying Markov chain. In

that case, the probability $\mathcal{P}_N(\boldsymbol{\mu})$ would be a product of multinomial distributions

$$\mathcal{P}_N(\boldsymbol{\mu}) = (N+1)! \prod_{V \in \mathcal{V}} \frac{p_V^{N_V}}{N_V!} N! \prod_{T \in \mathcal{T}} \frac{p_T^{N_T}}{N_T!} N! \prod_{\lambda \in \mathcal{L}} \frac{p_\lambda^{N_\lambda}}{N_\lambda!}, \quad (41)$$

where the multiplicities N_V , N_T and N_λ are integers which satisfy the constraints (24).

Equation (41) greatly simplifies for N large. By using Stirling's approximation for the factorials and explicitly taking into account the constraints (24), we have the following asymptotic equality for the associated probability density.

$$\begin{aligned} \mathcal{P}_N(N\boldsymbol{\nu}) \simeq \exp[N\omega(\boldsymbol{\nu})] \delta \left(\sum_{V \in \mathcal{V}} N\nu_V - N \right) \\ \times \delta \left(\sum_{T \in \mathcal{T}} N\nu_T - N \right) \delta \left(\sum_{\lambda \in \mathcal{L}} N\nu_\lambda - N \right), \end{aligned} \quad (42)$$

where

$$\omega(\boldsymbol{\nu}) = \sum_{\alpha \in \mathcal{H}} \nu_\alpha \log \left(\frac{p_\alpha}{\nu_\alpha} \right) \quad (43)$$

and $\mathbf{p}^T = (\dots p_V \dots; \dots p_T \dots; \dots p_\lambda \dots)$ is a vector collecting the success probabilities (38), (39) and (40). To find E_0 it remains to calculate the integral of equation (37) over the frequencies $\boldsymbol{\nu}$. The integration can be performed by steepest descent as detailed in the following section. The result is that E_0 is the solution of

$$\sum_{V \in \mathcal{V}} \frac{p_V}{-E_0 + V} = \frac{1}{(\sum_{T \in \mathcal{T}} p_T T) (\sum_{\lambda \in \mathcal{L}} p_\lambda \lambda)}, \quad E_0 \leq V_{\min}. \quad (44)$$

For $\hat{V} = 0$, i.e. $\mathcal{V} = \{0\}$ and $p_{V=0} = 1$, equation (44) can be solved analytically and we obtain the value

$$E_0^{(0)} = - \left(\sum_{T \in \mathcal{T}} p_T T \right) \left(\sum_{\lambda \in \mathcal{L}} p_\lambda \lambda \right) \quad (45)$$

for the ground-state energy of the hopping operator \hat{K} . For $\hat{V} \neq 0$ equation (44) is straightforwardly solved numerically by bisection method.

The uncorrelated multinomial probability density considered here has the great advantage that cumulants of any order are included in the determination of E_0 . This eliminates the artifacts obtained by truncating the cumulant expansion (32), namely the wrong behavior of E_0 at large interaction energies and its complex value in the case of fermions. In fact, for large interaction strength equation (44) admits the asymptotic finite solution

$$E_0 = V_{\min} + p_{V_{\min}} E_0^{(0)}. \quad (46)$$

Moreover, for intermediate interaction strengths the solution of equation (44) varies monotonously between the two limits (45) and (46), as expected. In the case of fermions, the $\hat{V} = 0$ solution (45) is real and negative, as must be. In fact, among the links $\lambda_{\mathbf{n}, \mathbf{n}'} \neq 0$ those with a positive sign are the majority so that

$$\sum_{\lambda \in \mathcal{L}} p_\lambda \lambda = \Sigma_{\lambda=1}^{(1)} - \Sigma_{\lambda=-1}^{(1)} > 0. \quad (47)$$

The main drawback of the present approach has already been mentioned. The correlations among the multiplicities $\boldsymbol{\mu}$ are neglected so that the comparison of E_0 , solution of equation (44), with the ground-state energy of a system of particles can be quantitatively poor. Whenever these correlations are absent, as in the case of the uniformly fully connected models and of the random potential systems considered in [12], the approach provides exact results together with the possibility to study the appearance of quantum phase transitions in the thermodynamic limit.

3. Multinomial perturbative scheme

Here we propose a probabilistic perturbative scheme with the aim of merging the merits of the multinomial probability density of section 2.2 with the statistical details provided by the cumulant expansion of section 2.1. Basically the idea is as follows. We consider an asymptotic probability density which has the same structure of a multinomial density but with parameters p_α , $\alpha \in \mathcal{H}$, that are functions of the frequencies $\boldsymbol{\nu}$,

$$\omega(\boldsymbol{\nu}) = \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathcal{P}_N(N\boldsymbol{\nu}) = \sum_{\alpha \in \mathcal{H}} \nu_\alpha \log \left(\frac{p_\alpha(\boldsymbol{\nu})}{\nu_\alpha} \right), \quad (48)$$

and write these functions as power series of the form

$$p_\alpha(\boldsymbol{\nu}) = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{\beta_1 \in \mathcal{H}} \cdots \sum_{\beta_n \in \mathcal{H}} p_{\alpha\beta_1 \dots \beta_n}^{(n+1)} \left(\nu_{\beta_1} - p_{\beta_1}^{(1)} \right) \cdots \left(\nu_{\beta_n} - p_{\beta_n}^{(1)} \right). \quad (49)$$

The scalars $p_{\alpha_1 \dots \alpha_k}^{(k)}$, with $\alpha_i \in \mathcal{H}$ for $i = 1, \dots, k$, in a compact notation $\mathbf{p}^{(k)}$, are the perturbative parameters at order k . Note that we identify the perturbative order k with the rank of the tensor $\mathbf{p}^{(k)}$ not the index n of the series (49), i.e. $k = n + 1 = 1, 2, \dots$. At the lowest perturbative order $k = 1$ we have $p_\alpha(\boldsymbol{\nu}) = p_\alpha^{(1)}$, $\alpha \in \mathcal{H}$, constant as in the strict multinomial case.

We determine the perturbative parameters $\mathbf{p}^{(k)}$ as follows. First, we assume that for $k \geq 2$ they are symmetric under the exchange of any two of their indices $\alpha_1, \dots, \alpha_k$

$$p_{\dots \alpha_i \dots \alpha_j \dots}^{(k)} = p_{\dots \alpha_j \dots \alpha_i \dots}^{(k)}. \quad (50)$$

Second, in order to maintain as much as possible the structure of a multinomial, we ask that, for any value of $\boldsymbol{\nu}$, the functions $p_\alpha(\boldsymbol{\nu})$ are normalized in each set $\mathcal{V}, \mathcal{I}, \mathcal{L}$, i.e.

$$\sum_{\alpha \in \mathcal{A}} p_\alpha(\boldsymbol{\nu}) = 1, \quad \mathcal{A} = \mathcal{V}, \mathcal{I}, \mathcal{L}. \quad (51)$$

Third, we require that the asymptotic rescaled cumulants of order k evaluated from (48–49), hereafter indicated by $\langle\langle \nu_{\alpha_1} \dots \nu_{\alpha_k} \rangle\rangle$, coincide with those effectively owned by the system, $\boldsymbol{\Sigma}^{(k)}$,

$$\langle\langle \nu_{\alpha_1} \dots \nu_{\alpha_k} \rangle\rangle(\mathbf{p}^{(1)}, \mathbf{p}^{(2)}, \dots, \mathbf{p}^{(k)}) = \boldsymbol{\Sigma}_{\alpha_1 \dots \alpha_k}^{(k)}. \quad (52)$$

In this expression we have anticipated that $\langle\langle \nu_{\alpha_1} \dots \nu_{\alpha_k} \rangle\rangle$ depends only on the parameters $\mathbf{p}^{(j)}$ with $j \leq k$. This property implies that we can first find $\mathbf{p}^{(1)}$ by solving the system of $|\mathcal{H}|$ equations

$$\langle\langle \nu_{\alpha_1} \rangle\rangle(\mathbf{p}^{(1)}) = \boldsymbol{\Sigma}_{\alpha_1}^{(1)}, \quad \alpha_1 \in \mathcal{H}, \quad (53)$$

next find $\mathbf{p}^{(2)}$ by solving the system of $|\mathcal{H}|^2$ equations

$$\langle\langle \nu_{\alpha_1} \nu_{\alpha_2} \rangle\rangle(\mathbf{p}^{(1)}, \mathbf{p}^{(2)}) = \Sigma_{\alpha_1 \alpha_2}^{(2)}, \quad \alpha_1, \alpha_2 \in \mathcal{H}, \quad (54)$$

next find $\mathbf{p}^{(3)}$ by solving the system of $|\mathcal{H}|^3$ equations

$$\langle\langle \nu_{\alpha_1} \nu_{\alpha_2} \nu_{\alpha_3} \rangle\rangle(\mathbf{p}^{(1)}, \mathbf{p}^{(2)}, \mathbf{p}^{(3)}) = \Sigma_{\alpha_1 \alpha_2 \alpha_3}^{(3)}, \quad \alpha_1, \alpha_2, \alpha_3 \in \mathcal{H}, \quad (55)$$

and so on up to a chosen maximum order k_{\max} which corresponds to truncate the series (49) at the term $n = k_{\max} - 1$ included.

To be precise, the probability density (48) is well defined only if each $p_\alpha(\boldsymbol{\nu})$, $\alpha \in \mathcal{H}$, is a non negative function of the vector $\boldsymbol{\nu}$ with components $\nu_\alpha \geq 0$ varying in the unit simplex $\sum_{\alpha \in \mathcal{A}} \nu_\alpha = 1$ for $\mathcal{A} = \mathcal{V}, \mathcal{T}, \mathcal{L}$. Actually, we will content ourselves that the positivity condition for $\mathbf{p}(\boldsymbol{\nu})$ is satisfied in a small region around the frequency $\boldsymbol{\nu}$ which gives an exponentially leading contribution to the integral (37). *A priori* we don't know whether this condition can be met by introducing the perturbative parameters $\mathbf{p}^{(k)}$ as described above. We will then proceed heuristically. Whenever an effective solution of equation (37) can be found by using the probability density (48–49), that will be the signal that the proposed perturbative scheme is meaningful.

In the following subsections, we will first evaluate, up to the third order, the asymptotic rescaled cumulants associated to the probability density (48–49), then we will discuss the solution of the systems of equations (53), (54) and (55). Finally, we will make some comments on the higher perturbative orders.

3.1. Evaluation of the asymptotic rescaled cumulants

The definition of the asymptotic rescaled cumulants of order $k = 1, 2, \dots$ is

$$\langle\langle \nu_{\alpha_1} \dots \nu_{\alpha_k} \rangle\rangle = \lim_{N \rightarrow \infty} \frac{1}{N} \langle N \nu_{\alpha_1} \dots N \nu_{\alpha_k} \rangle_N^{(c)}, \quad \alpha_1, \dots, \alpha_k \in \mathcal{H}, \quad (56)$$

where $\langle N \nu_{\alpha_1} \dots N \nu_{\alpha_k} \rangle_N^{(c)}$ are the cumulants, or connected correlation functions, of the multiplicities $N_\alpha = N \nu_\alpha$ sampled with respect to the N -jumps probability density $\mathcal{P}_N(N\boldsymbol{\nu})$. In turn, as standard, the cumulants are obtained from the generating function associated to $\mathcal{P}_N(N\boldsymbol{\nu})$

$$\langle N \nu_{\alpha_1} \dots N \nu_{\alpha_k} \rangle_N^{(c)} = \left. \frac{\partial^k \log Z_N(\mathbf{J})}{\partial J_{\alpha_1} \dots \partial J_{\alpha_k}} \right|_{\mathbf{J}=\mathbf{0}}, \quad (57)$$

$$Z_N(\mathbf{J}) = \int d(N\boldsymbol{\nu}) \mathcal{P}_N(N\boldsymbol{\nu}) e^{(\mathbf{J}, N\boldsymbol{\nu})}. \quad (58)$$

Assuming that for $N \rightarrow \infty$ the logarithm of $\mathcal{P}_N(N\boldsymbol{\nu})$ is given by (48–49), for N large we have, up to an inessential constant,

$$Z_N(\mathbf{J}) = \int d(N\boldsymbol{\nu}) e^{N\omega(\boldsymbol{\nu}) + (\mathbf{J}, N\boldsymbol{\nu})} \prod_{\mathcal{A}=\mathcal{V}, \mathcal{T}, \mathcal{L}} \delta \left(\sum_{\alpha \in \mathcal{A}} N \nu_\alpha - N \right), \quad (59)$$

where we have explicitly taken into account the fact that the multiplicities of each set $\mathcal{V}, \mathcal{T}, \mathcal{L}$ must sum to N . Using the Fourier integral representation of the Dirac δ , we

rewrite the generating function, up to a constant, as

$$Z_N(\mathbf{J}) = \int \prod_{\mathcal{A}=\mathcal{V},\mathcal{T},\mathcal{L}} dk_{\mathcal{A}} \prod_{\alpha \in \mathcal{H}} d\nu_{\alpha} e^{N\phi(\boldsymbol{\nu}, \mathbf{k}, \mathbf{J})}, \quad (60)$$

where

$$\begin{aligned} \phi(\boldsymbol{\nu}, \mathbf{k}, \mathbf{J}) &= \sum_{\alpha \in \mathcal{H}} \nu_{\alpha} \left[\log \left(\frac{p_{\alpha}(\boldsymbol{\nu})}{\nu_{\alpha}} \right) + J_{\alpha} \right] \\ &+ \sum_{\mathcal{A}=\mathcal{V},\mathcal{T},\mathcal{L}} ik_{\mathcal{A}} \left(\sum_{\alpha \in \mathcal{A}} \nu_{\alpha} - 1 \right). \end{aligned} \quad (61)$$

In the above expressions we used the compact notation $\mathbf{k} = (k_{\mathcal{V}}, k_{\mathcal{T}}, k_{\mathcal{L}})$ and $\mathbf{J} = (\dots J_V \dots; \dots J_T \dots; \dots J_{\lambda} \dots)$ with $V \in \mathcal{V}$, $T \in \mathcal{T}$ and $\lambda \in \mathcal{L}$. Evaluating the integrals in equation (60) by Laplace's method, we get the N -large asymptotic logarithm equality

$$Z_N(\mathbf{J}) \simeq e^{N\phi_{\text{sp}}(\mathbf{J})}, \quad \phi_{\text{sp}}(\mathbf{J}) = \phi(\boldsymbol{\nu}^{\text{sp}}(\mathbf{J}), \mathbf{k}^{\text{sp}}(\mathbf{J}), \mathbf{J}), \quad (62)$$

where $(\boldsymbol{\nu}^{\text{sp}}, \mathbf{k}^{\text{sp}})$ is the saddle point (actually, the maximum point) of ϕ solution of the following system of equations

$$\frac{\partial \phi}{\partial \nu_{\alpha}} = 0, \quad \alpha \in \mathcal{H}, \quad (63)$$

$$\frac{\partial \phi}{\partial k_{\mathcal{A}}} = 0, \quad \mathcal{A} = \mathcal{V}, \mathcal{T}, \mathcal{L}. \quad (64)$$

By differentiating equation (61), the saddle-point equations can be written as

$$\log \left(\frac{\nu_{\alpha}^{\text{sp}}}{p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})} \right) = J_{\alpha} + \sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\beta}^{\text{sp}} + ik_{\mathcal{A}_{\alpha}}^{\text{sp}} - 1, \quad \alpha \in \mathcal{H}, \quad (65)$$

$$\sum_{\alpha \in \mathcal{A}} \nu_{\alpha}^{\text{sp}} = 1, \quad \mathcal{A} = \mathcal{V}, \mathcal{T}, \mathcal{L}, \quad (66)$$

where we have defined

$$\begin{aligned} p'_{\alpha\beta}(\boldsymbol{\nu}) &= \frac{\partial p_{\alpha}(\boldsymbol{\nu})}{\partial \nu_{\beta}} \\ &= p_{\alpha\beta}^{(2)} + \sum_{\gamma \in \mathcal{H}} p_{\alpha\beta\gamma}^{(3)} (\nu_{\gamma} - p_{\gamma}^{(1)}) + \dots \end{aligned} \quad (67)$$

and

$$\mathcal{A}_{\alpha} = \begin{cases} \mathcal{V}, & \alpha \in \mathcal{V}, \\ \mathcal{T}, & \alpha \in \mathcal{T}, \\ \mathcal{L}, & \alpha \in \mathcal{L}. \end{cases} \quad (68)$$

Note that, due to the properties assumed for the perturbative parameters, $p'_{\alpha\beta}(\boldsymbol{\nu})$ is symmetric under the exchange of the indices α, β . From equation (65) we find

$$\nu_{\alpha}^{\text{sp}} = p_{\alpha}(\boldsymbol{\nu}^{\text{sp}}) e^{J_{\alpha} + \sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\beta}^{\text{sp}} + ik_{\mathcal{A}_{\alpha}}^{\text{sp}} - 1}, \quad \alpha \in \mathcal{H}, \quad (69)$$

which inserted into equation (66) provides

$$e^{ik_{\mathcal{A}}^{\text{sp}}-1} = \frac{1}{\sum_{\alpha \in \mathcal{A}} p_{\alpha}(\boldsymbol{\nu}^{\text{sp}}) e^{J_{\alpha} + \sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\beta}^{\text{sp}}}}, \quad \mathcal{A} = \mathcal{V}, \mathcal{T}, \mathcal{L}. \quad (70)$$

By using equation (70) we conclude that the saddle-point frequencies (69) are the solution of the system of nonlinear equations

$$\nu_{\alpha}^{\text{sp}} = \frac{p_{\alpha}(\boldsymbol{\nu}^{\text{sp}}) e^{J_{\alpha} + \sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\beta}^{\text{sp}}}}{\sum_{\alpha' \in \mathcal{A}_{\alpha}} p_{\alpha'}(\boldsymbol{\nu}^{\text{sp}}) e^{J_{\alpha'} + \sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha'\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\beta}^{\text{sp}}}}, \quad \alpha \in \mathcal{H}. \quad (71)$$

Note that both $\boldsymbol{\nu}^{\text{sp}}$ and \mathbf{k}^{sp} are functions of the source \mathbf{J} . It follows that the function $\phi(\boldsymbol{\nu}, \mathbf{k}, \mathbf{J})$ evaluated at the saddle point $(\boldsymbol{\nu}^{\text{sp}}(\mathbf{J}), \mathbf{k}^{\text{sp}}(\mathbf{J}))$ is

$$\begin{aligned} \phi_{\text{sp}}(\mathbf{J}) &= \sum_{\alpha \in \mathcal{H}} \nu_{\alpha}^{\text{sp}} \left[\log \left(\frac{p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})}{\nu_{\alpha}^{\text{sp}}} \right) + J_{\alpha} \right] \\ &= \sum_{\mathcal{A}=\mathcal{V}, \mathcal{T}, \mathcal{L}} \log \left(\sum_{\alpha \in \mathcal{A}} p_{\alpha}(\boldsymbol{\nu}^{\text{sp}}) e^{J_{\alpha} + \sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\beta}^{\text{sp}}} \right) \\ &\quad - \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\alpha}^{\text{sp}} \nu_{\beta}^{\text{sp}}. \end{aligned} \quad (72)$$

We are now ready to evaluate the asymptotic rescaled cumulants associated to the probability density (48–49) by using the formula

$$\langle\langle \nu_{\alpha_1} \dots \nu_{\alpha_k} \rangle\rangle = \left. \frac{\partial^k \phi_{\text{sp}}(\mathbf{J})}{\partial J_{\alpha_1} \dots \partial J_{\alpha_k}} \right|_{\mathbf{J}=\mathbf{0}}. \quad (73)$$

3.2. Derivatives of $\phi_{\text{sp}}(\mathbf{J})$

In this subsection we supply the derivatives of $\phi_{\text{sp}}(\mathbf{J})$ with respect to the source \mathbf{J} up to the third order. Note that ϕ_{sp} depends on \mathbf{J} explicitly and through the saddle-point frequencies.

The first order derivative of $\phi_{\text{sp}}(\mathbf{J})$ with respect to the source component J_{α_1} , $\alpha_1 \in \mathcal{H}$, is

$$\begin{aligned} \frac{\partial \phi_{\text{sp}}(\mathbf{J})}{\partial J_{\alpha_1}} &= \sum_{\mathcal{A}=\mathcal{V}, \mathcal{T}, \mathcal{L}} \sum_{\alpha \in \mathcal{A}} \nu_{\alpha}^{\text{sp}} \left(\frac{1}{p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})} \frac{\partial p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})}{\partial J_{\alpha_1}} + \delta_{\alpha\alpha_1} \right) \\ &\quad - \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \frac{\partial \nu_{\alpha}^{\text{sp}}}{\partial J_{\alpha_1}} \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\beta}^{\text{sp}}. \end{aligned}$$

Since

$$\frac{\partial p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})}{\partial J_{\alpha_1}} = \sum_{\beta \in \mathcal{H}} p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}}) \frac{\partial \nu_{\beta}^{\text{sp}}}{\partial J_{\alpha_1}} \quad (74)$$

and $p'_{\alpha\beta}$ is symmetric, the previous expression reduces to

$$\frac{\partial \phi_{\text{sp}}(\mathbf{J})}{\partial J_{\alpha_1}} = \nu_{\alpha_1}^{\text{sp}}. \quad (75)$$

The second order derivative of $\phi_{\text{sp}}(\mathbf{J})$ with respect to the source components J_{α_1} and J_{α_2} , $\alpha_1, \alpha_2 \in \mathcal{H}$, is

$$\begin{aligned} \frac{\partial^2 \phi_{\text{sp}}(\mathbf{J})}{\partial J_{\alpha_1} \partial J_{\alpha_2}} &= \frac{\partial \nu_{\alpha_1}^{\text{sp}}}{\partial J_{\alpha_2}} \\ &= \nu_{\alpha_1}^{\text{sp}} \delta_{\alpha_1 \alpha_2} - \nu_{\alpha_1}^{\text{sp}} \nu_{\alpha_2}^{\text{sp}} \chi_{\alpha_1 \alpha_2} \\ &\quad + \sum_{\alpha \in \mathcal{H}} (\nu_{\alpha_1}^{\text{sp}} \delta_{\alpha_1 \alpha} - \nu_{\alpha_1}^{\text{sp}} \nu_{\alpha}^{\text{sp}} \chi_{\alpha_1 \alpha}) \\ &\quad \times \left[\frac{1}{p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})} \frac{\partial p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})}{\partial J_{\alpha_2}} + \sum_{\beta \in \mathcal{H}} \frac{\partial}{\partial J_{\alpha_2}} \left(\frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\beta}^{\text{sp}} \right) \right], \end{aligned}$$

where

$$\chi_{\alpha\beta} = \chi_{\beta\alpha} = \begin{cases} 1, & \alpha \in \mathcal{A}_{\beta}, \\ 0, & \text{otherwise.} \end{cases} \quad (76)$$

Recalling the definition (67) and using equation (74), we find

$$\begin{aligned} \frac{\partial^2 \phi_{\text{sp}}(\mathbf{J})}{\partial J_{\alpha_1} \partial J_{\alpha_2}} &= \nu_{\alpha_1}^{\text{sp}} \delta_{\alpha_1 \alpha_2} - \nu_{\alpha_1}^{\text{sp}} \nu_{\alpha_2}^{\text{sp}} \chi_{\alpha_1 \alpha_2} \\ &\quad + \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} (\nu_{\alpha_1}^{\text{sp}} \delta_{\alpha_1 \alpha} - \nu_{\alpha_1}^{\text{sp}} \nu_{\alpha}^{\text{sp}} \chi_{\alpha_1 \alpha}) M_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}}) \frac{\partial \nu_{\beta}^{\text{sp}}}{\partial J_{\alpha_2}}, \end{aligned} \quad (77)$$

where we have introduced the symmetric matrix

$$M_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}}) = \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})} + \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} + \sum_{\gamma \in \mathcal{H}} \left(\frac{p''_{\alpha\beta\gamma}(\boldsymbol{\nu}^{\text{sp}})}{p_{\gamma}(\boldsymbol{\nu}^{\text{sp}})} - \frac{p'_{\alpha\gamma}(\boldsymbol{\nu}^{\text{sp}}) p'_{\gamma\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\gamma}(\boldsymbol{\nu}^{\text{sp}})^2} \right) \nu_{\gamma}^{\text{sp}} \quad (78)$$

and defined

$$\begin{aligned} p''_{\alpha\beta\gamma}(\boldsymbol{\nu}) &= \frac{\partial^2 p_{\alpha}(\boldsymbol{\nu})}{\partial \nu_{\beta} \partial \nu_{\gamma}} \\ &= p_{\alpha\beta\gamma}^{(3)} + \sum_{\delta \in \mathcal{H}} p_{\alpha\beta\gamma\delta}^{(4)} (\nu_{\delta} - p_{\delta}^{(1)}) + \dots \end{aligned} \quad (79)$$

Note that, due to the properties assumed for the perturbative parameters, $p''_{\alpha\beta\gamma}(\boldsymbol{\nu})$ is symmetric under the exchange of any two of its indices α, β, γ .

The third order derivative of $\phi_{\text{sp}}(\mathbf{J})$ with respect to the source components J_{α_1} , J_{α_2} and J_{α_3} , $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{H}$, is

$$\begin{aligned} \frac{\partial^3 \phi_{\text{sp}}(\mathbf{J})}{\partial J_{\alpha_1} \partial J_{\alpha_2} \partial J_{\alpha_3}} &= \frac{\partial^2 \nu_{\alpha_1}^{\text{sp}}}{\partial J_{\alpha_2} \partial J_{\alpha_3}} \\ &= \frac{\partial \nu_{\alpha_1}^{\text{sp}}}{\partial J_{\alpha_3}} \delta_{\alpha_1 \alpha_2} - \left(\frac{\partial \nu_{\alpha_1}^{\text{sp}}}{\partial J_{\alpha_3}} \nu_{\alpha_2}^{\text{sp}} + \nu_{\alpha_1}^{\text{sp}} \frac{\partial \nu_{\alpha_2}^{\text{sp}}}{\partial J_{\alpha_3}} \right) \chi_{\alpha_1 \alpha_2} \\ &\quad + \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \left(\frac{\partial \nu_{\alpha_1}^{\text{sp}}}{\partial J_{\alpha_3}} \delta_{\alpha_1 \alpha} - \left(\frac{\partial \nu_{\alpha_1}^{\text{sp}}}{\partial J_{\alpha_3}} \nu_{\alpha}^{\text{sp}} + \nu_{\alpha_1}^{\text{sp}} \frac{\partial \nu_{\alpha}^{\text{sp}}}{\partial J_{\alpha_3}} \right) \chi_{\alpha_1 \alpha} \right) M_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}}) \frac{\partial \nu_{\beta}^{\text{sp}}}{\partial J_{\alpha_2}} \\ &\quad + \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} (\nu_{\alpha_1}^{\text{sp}} \delta_{\alpha_1 \alpha} - \nu_{\alpha_1}^{\text{sp}} \nu_{\alpha}^{\text{sp}} \chi_{\alpha_1 \alpha}) \\ &\quad \times \left(M_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}}) \frac{\partial^2 \nu_{\beta}^{\text{sp}}}{\partial J_{\alpha_2} \partial J_{\alpha_3}} + \frac{\partial M_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{\partial J_{\alpha_3}} \frac{\partial \nu_{\beta}^{\text{sp}}}{\partial J_{\alpha_2}} \right), \end{aligned} \quad (80)$$

with

$$\begin{aligned}
\frac{\partial M_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{\partial J_{\alpha_3}} = & \sum_{\gamma \in \mathcal{H}} \left[\frac{p''_{\alpha\beta\gamma}(\boldsymbol{\nu}^{\text{sp}})}{p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})} + \frac{p''_{\alpha\beta\gamma}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} + \frac{p''_{\alpha\beta\gamma}(\boldsymbol{\nu}^{\text{sp}})}{p_{\gamma}(\boldsymbol{\nu}^{\text{sp}})} \right. \\
& - \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})p'_{\alpha\gamma}(\boldsymbol{\nu}^{\text{sp}})}{p_{\alpha}(\boldsymbol{\nu}^{\text{sp}})^2} - \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})p'_{\beta\gamma}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})^2} - \frac{p'_{\alpha\gamma}(\boldsymbol{\nu}^{\text{sp}})p'_{\gamma\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\gamma}(\boldsymbol{\nu}^{\text{sp}})^2} \\
& + \sum_{\delta \in \mathcal{H}} \left(\frac{p'''_{\alpha\beta\gamma\delta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\delta}(\boldsymbol{\nu}^{\text{sp}})} - \frac{p''_{\alpha\beta\delta}(\boldsymbol{\nu}^{\text{sp}})p'_{\delta\gamma}(\boldsymbol{\nu}^{\text{sp}})}{p_{\delta}(\boldsymbol{\nu}^{\text{sp}})^2} \right. \\
& - \frac{p''_{\alpha\delta\gamma}(\boldsymbol{\nu}^{\text{sp}})p'_{\delta\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\delta}(\boldsymbol{\nu}^{\text{sp}})^2} - \frac{p''_{\delta\beta\gamma}(\boldsymbol{\nu}^{\text{sp}})p'_{\delta\alpha}(\boldsymbol{\nu}^{\text{sp}})}{p_{\delta}(\boldsymbol{\nu}^{\text{sp}})^2} \\
& \left. \left. + 2 \frac{p'_{\alpha\delta}(\boldsymbol{\nu}^{\text{sp}})p'_{\beta\delta}(\boldsymbol{\nu}^{\text{sp}})p'_{\gamma\delta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\delta}(\boldsymbol{\nu}^{\text{sp}})^3} \right) \nu_{\delta}^{\text{sp}} \right] \frac{\partial \nu_{\gamma}^{\text{sp}}}{\partial J_{\alpha_3}}, \tag{81}
\end{aligned}$$

where we have defined

$$\begin{aligned}
p'''_{\alpha\beta\gamma\delta}(\boldsymbol{\nu}) &= \frac{\partial^3 p_{\alpha}(\boldsymbol{\nu})}{\partial \nu_{\beta} \partial \nu_{\gamma} \partial \nu_{\delta}} \\
&= p_{\alpha\beta\gamma\delta}^{(4)} + \sum_{\varepsilon \in \mathcal{H}} p_{\alpha\beta\gamma\delta\varepsilon}^{(5)} (\nu_{\varepsilon} - p_{\varepsilon}^{(1)}) + \dots
\end{aligned} \tag{82}$$

3.3. Equations for the perturbative parameters: first order

The perturbative parameters $\mathbf{p}^{(1)}$ are determined by the system of equations (53) which, according to (73) and (75), reads

$$\nu_{\alpha_1}^{\text{sp}}|_{\mathbf{J}=\mathbf{0}} = \Sigma_{\alpha_1}^{(1)}, \quad \alpha_1 \in \mathcal{H}. \tag{83}$$

Using equation (71), we explicitly have

$$\begin{aligned}
\frac{p_{\alpha_1}(\boldsymbol{\Sigma}^{(1)}) e^{\sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha_1\beta}(\boldsymbol{\Sigma}^{(1)})}{p_{\beta}(\boldsymbol{\Sigma}^{(1)})} \Sigma_{\beta}^{(1)}}}{\sum_{\alpha \in \mathcal{A}_{\alpha_1}} p_{\alpha}(\boldsymbol{\Sigma}^{(1)}) e^{\sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha\beta}(\boldsymbol{\Sigma}^{(1)})}{p_{\beta}(\boldsymbol{\Sigma}^{(1)})} \Sigma_{\beta}^{(1)}}} &= \Sigma_{\alpha_1}^{(1)}, \quad \alpha_1 \in \mathcal{H}.
\end{aligned} \tag{84}$$

A solution of this system of nonlinear equations is

$$p_{\alpha_1}^{(1)} = \Sigma_{\alpha_1}^{(1)}, \quad \alpha_1 \in \mathcal{H}. \tag{85}$$

To prove it, we start to observe that if $\mathbf{p}^{(1)} = \boldsymbol{\Sigma}^{(1)}$ then for any $\alpha, \beta \in \mathcal{H}$

$$p_{\alpha}(\boldsymbol{\Sigma}^{(1)}) = p_{\alpha}^{(1)}, \tag{86}$$

and

$$p'_{\alpha\beta}(\boldsymbol{\Sigma}^{(1)}) = p_{\alpha\beta}^{(2)}. \tag{87}$$

The position $\mathbf{p}^{(1)} = \boldsymbol{\Sigma}^{(1)}$ has another consequence which is pivotal also in determining the perturbative parameters of higher order. In fact, due to the constraints (51), the analogous normalization conditions valid for $\boldsymbol{\Sigma}^{(1)}$ and the symmetry properties (50), we have

$$\sum_{\alpha_i \in \mathcal{A}} p_{\alpha_1 \dots \alpha_k}^{(k)} = 0, \quad k \geq 2, \quad 1 \leq i \leq k, \quad \mathcal{A} = \mathcal{V}, \mathcal{I}, \mathcal{L}. \tag{88}$$

In the present case, the above sum rules allow to write $\sum_{\beta \in \mathcal{H}} p_{\alpha\beta}^{(2)} = 0$, which, together with (86) and (87), permits to reduce the l.h.s. of equation (84) to $\Sigma_{\alpha_1}^{(1)}$.

Equation (85) is not the unique solution of the system (84). However, besides being the natural solution for which the perturbed probability density (48-49) reduces, for $k_{\max} = 1$, to the uncorrelated multinomial (43), it allows us to establish the sum rules (88). We shall show that these, in turn, entail the asymptotic rescaled cumulants $\langle\langle \nu_{\alpha_1} \dots \nu_{\alpha_k} \rangle\rangle$ of arbitrary order k to depend only by the parameters $\mathbf{p}^{(j)}$ with $j \leq k$. In this way, the parameters $\mathbf{p}^{(1)}, \mathbf{p}^{(2)}, \mathbf{p}^{(3)}, \dots$ we find do not depend on the order k_{\max} at which we decide to truncate the series (49).

3.4. Equations for the perturbative parameters: second order

The perturbative parameters $\mathbf{p}^{(2)}$ are determined by the system of equations (54) which, according to (73) and (77), reads

$$\Sigma_{\alpha_1\alpha_2}^{(2,0)} + \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1\alpha}^{(2,0)} \left[\frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\alpha}^{(1)}} + \frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\beta}^{(1)}} + \sum_{\gamma \in \mathcal{H}} \left(\frac{p_{\alpha\beta\gamma}^{(3)}}{\Sigma_{\gamma}^{(1)}} - \frac{p_{\alpha\gamma}^{(2)} p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)^2}} \right) \Sigma_{\gamma}^{(1)} \right] \Sigma_{\beta\alpha_2}^{(2)} = \Sigma_{\alpha_1\alpha_2}^{(2)}.$$

In writing this expression we have used the results (83), (85), (86) and (87) of the previous section as well as the fact that, since $\mathbf{p}^{(1)} = \Sigma^{(1)}$, for any $\alpha, \beta, \gamma \in \mathcal{H}$ it is

$$p''_{\alpha\beta\gamma}(\Sigma^{(1)}) = p_{\alpha\beta\gamma}^{(3)}. \quad (89)$$

We have also defined

$$\Sigma_{\alpha\beta}^{(2,0)} = \Sigma_{\alpha}^{(1)} \delta_{\alpha\beta} - \Sigma_{\alpha}^{(1)} \Sigma_{\beta}^{(1)} \chi_{\alpha\beta}, \quad \alpha, \beta \in \mathcal{H}, \quad (90)$$

$\Sigma^{(2,0)}$ in a compact notation, which is the second cumulant of an uncorrelated multinomial probability density with parameters $\Sigma^{(1)}$. According to equation (88), we have $\sum_{\gamma \in \mathcal{H}} p_{\alpha\beta\gamma}^{(3)} = 0$ so that the above system of equations becomes

$$\sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1\alpha}^{(2,0)} \left[\frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\alpha}^{(1)}} + \frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\beta}^{(1)}} - \sum_{\gamma \in \mathcal{H}} \frac{p_{\alpha\gamma}^{(2)} p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)}} \right] \Sigma_{\beta\alpha_2}^{(2)} = \Sigma_{\alpha_1\alpha_2}^{(2)} - \Sigma_{\alpha_1\alpha_2}^{(2,0)}. \quad (91)$$

For $\alpha_1, \alpha_2 \in \mathcal{H}$, this is a nonlinear system of m^2 , $m = |\mathcal{H}|$, equations. On the other hand, $\mathbf{p}^{(2)}$, recall that it satisfies the sum rules (88), has \hat{m}^2 , $\hat{m} = (|\mathcal{V}| - 1) + (|\mathcal{T}| - 1) + (|\mathcal{L}| - 1)$, independent components. Therefore, the system of equations (91) is overdetermined and we have to lower its dimension to find $\mathbf{p}^{(2)}$. Note that also the $m \times m$ matrices $\Sigma^{(2)}$ and $\Sigma^{(2,0)}$ are singular and their rank is \hat{m}^2 .

Let us introduce the \hat{m} -dimensional index space $\hat{\mathcal{H}} = \mathcal{H} \setminus \{V_*, T_*, \lambda_*\}$, where V_* , T_* and λ_* are three arbitrarily chosen elements of the sets \mathcal{V} , \mathcal{T} and \mathcal{L} , respectively. Let $\hat{\mathbf{p}}^{(1)}$ be the vector with components $\hat{p}_{\alpha}^{(1)} = p_{\alpha}^{(1)}$, $\alpha \in \hat{\mathcal{H}}$, and $\hat{\mathbf{p}}^{(2)}$ the matrix with components $\hat{p}_{\alpha\beta}^{(2)} = p_{\alpha\beta}^{(2)}$, $\alpha, \beta \in \hat{\mathcal{H}}$. Now consider the \hat{m}^2 equations (91) with $\alpha_1, \alpha_2 \in \hat{\mathcal{H}}$. Observing that $\sum_{\alpha \in \mathcal{H}} = \sum_{\alpha \in \hat{\mathcal{H}}} + \sum_{\alpha \in \{V_*, T_*, \lambda_*\}}$ and using the sum rules (88), we rewrite the first term in the l.h.s. of equation (91) as

$$\sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1\alpha}^{(2,0)} \frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\alpha}^{(1)}} \Sigma_{\beta\alpha_2}^{(2)} = \sum_{\alpha \in \hat{\mathcal{H}}} \sum_{\beta \in \hat{\mathcal{H}}} \left(\frac{\Sigma_{\alpha_1\alpha}^{(2,0)}}{\Sigma_{\alpha}^{(1)}} - \frac{\Sigma_{\alpha_1\alpha_*}^{(2,0)}}{\Sigma_{\alpha_*}^{(1)}} \right) \hat{p}_{\alpha\beta}^{(2)} \left(\Sigma_{\beta\alpha_2}^{(2)} - \Sigma_{\beta_*\alpha_2}^{(2)} \right),$$

where α_* , β_* are the components eliminated from the sets \mathcal{A}_α , \mathcal{A}_β , i.e.

$$\alpha_* = \begin{cases} V_*, & \alpha \in \mathcal{V}, \\ T_*, & \alpha \in \mathcal{T}, \\ \lambda_*, & \alpha \in \mathcal{L}. \end{cases} \quad (92)$$

Similarly, the second term in the l.h.s. of (91) becomes

$$\sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1 \alpha}^{(2,0)} \frac{p_{\alpha\beta}^{(2)}}{\Sigma_\beta^{(1)}} \Sigma_{\beta\alpha_2}^{(2)} = \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} (\Sigma_{\alpha_1 \alpha}^{(2,0)} - \Sigma_{\alpha_1 \alpha_*}^{(2,0)}) \hat{p}_{\alpha\beta}^{(2)} \left(\frac{\Sigma_{\beta\alpha_2}^{(2)}}{\Sigma_\beta^{(1)}} - \frac{\Sigma_{\beta_*\alpha_2}^{(2)}}{\Sigma_{\beta_*}^{(1)}} \right),$$

whereas the third term gives

$$\begin{aligned} & \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \sum_{\gamma \in \mathcal{H}} \Sigma_{\alpha_1 \alpha}^{(2,0)} \frac{p_{\alpha\gamma}^{(2)} p_{\gamma\beta}^{(2)}}{\Sigma_\gamma^{(1)}} \Sigma_{\beta\alpha_2}^{(2)} \\ &= \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \sum_{\gamma \in \mathcal{H}} \sum_{\delta \in \mathcal{H}} (\Sigma_{\alpha_1 \alpha}^{(2,0)} - \Sigma_{\alpha_1 \alpha_*}^{(2,0)}) \hat{p}_{\alpha\gamma}^{(2)} \left(\frac{\delta_{\gamma\delta}}{\Sigma_\gamma^{(1)}} + \frac{\chi_{\gamma\delta}}{\Sigma_{\gamma_*}^{(1)}} \right) \hat{p}_{\delta\beta}^{(2)} (\Sigma_{\beta\alpha_2}^{(2)} - \Sigma_{\beta_*\alpha_2}^{(2)}). \end{aligned}$$

We conclude that $\hat{\mathbf{p}}^{(2)}$ is the solution of the nonlinear matrix equation

$$\bar{\Sigma}^{(2,0)} \hat{\mathbf{p}}^{(2)} \tilde{\Sigma}^{(2)} + \tilde{\Sigma}^{(2,0)} \hat{\mathbf{p}}^{(2)} \bar{\Sigma}^{(2)} - \tilde{\Sigma}^{(2,0)} \hat{\mathbf{p}}^{(2)} \Gamma \hat{\mathbf{p}}^{(2)} \tilde{\Sigma}^{(2)} = \hat{\Sigma}^{(2)} - \hat{\Sigma}^{(2,0)}. \quad (93)$$

where $\hat{\Sigma}^{(2)}$ and $\hat{\Sigma}^{(2,0)}$ are the reduced versions of the matrices $\Sigma^{(2)}$ and $\Sigma^{(2,0)}$ and we have introduced the matrices $\tilde{\Sigma}^{(2)}$, $\bar{\Sigma}^{(2)}$, $\tilde{\Sigma}^{(2,0)}$, $\bar{\Sigma}^{(2,0)}$ and Γ with components $\alpha, \beta \in \mathcal{H}$ given by

$$\tilde{\Sigma}_{\alpha\beta}^{(2)} = \Sigma_{\alpha\beta}^{(2)} - \Sigma_{\alpha_*\beta}^{(2)}, \quad (94)$$

$$\bar{\Sigma}_{\alpha\beta}^{(2)} = \frac{\Sigma_{\alpha\beta}^{(2)}}{\Sigma_\alpha^{(1)}} - \frac{\Sigma_{\alpha_*\beta}^{(2)}}{\Sigma_{\alpha_*}^{(1)}}, \quad (95)$$

$$\tilde{\Sigma}_{\alpha\beta}^{(2,0)} = \Sigma_{\alpha\beta}^{(2,0)} - \Sigma_{\alpha\beta_*}^{(2,0)}, \quad (96)$$

$$\bar{\Sigma}_{\alpha\beta}^{(2,0)} = \frac{\Sigma_{\alpha\beta}^{(2,0)}}{\Sigma_\beta^{(1)}} - \frac{\Sigma_{\alpha\beta_*}^{(2,0)}}{\Sigma_{\beta_*}^{(1)}}, \quad (97)$$

$$\Gamma_{\alpha\beta} = \frac{\delta_{\alpha\beta}}{\Sigma_\alpha^{(1)}} + \frac{\chi_{\alpha\beta}}{\Sigma_{\alpha_*}^{(1)}}. \quad (98)$$

Since $\tilde{\Sigma}^{(2)}$ and $\tilde{\Sigma}^{(2,0)}$ are nonsingular, equation (93) can be rewritten as

$$\begin{aligned} & \tilde{\Sigma}^{(2,0)-1} \bar{\Sigma}^{(2,0)} \hat{\mathbf{p}}^{(2)} + \hat{\mathbf{p}}^{(2)} \bar{\Sigma}^{(2)} \tilde{\Sigma}^{(2)-1} - \hat{\mathbf{p}}^{(2)} \Gamma \hat{\mathbf{p}}^{(2)} \\ &= \tilde{\Sigma}^{(2,0)-1} (\hat{\Sigma}^{(2)} - \hat{\Sigma}^{(2,0)}) \tilde{\Sigma}^{(2)-1}. \end{aligned} \quad (99)$$

This is a nonsymmetric algebraic Riccati equation (NARE), which can be solved numerically by matrix factorization (Schür method) [17] or iterative methods [18]. For details we refer to Appendix A. Once $\hat{\mathbf{p}}^{(2)}$ has been found, the complete set of second order perturbative parameters $\mathbf{p}^{(2)}$ is obtained using the sum rules (88) for $k = 2$.

3.5. Equations for the perturbative parameters: third order

The perturbative parameters $\mathbf{p}^{(3)}$ are determined by the system of equations (55) which, according to (73) and (80), reads

$$\begin{aligned}
\Sigma_{\alpha_1\alpha_2\alpha_3}^{(3,0)} &+ \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1\alpha\alpha_3}^{(3,0)} \left(\frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\alpha}^{(1)}} + \frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\beta}^{(1)}} + \sum_{\gamma \in \mathcal{H}} \left(\frac{p_{\alpha\beta\gamma}^{(3)}}{\Sigma_{\gamma}^{(1)}} - \frac{p_{\alpha\gamma}^{(2)}p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)^2}} \right) \Sigma_{\gamma}^{(1)} \right) \Sigma_{\beta\alpha_2}^{(2)} \\
&+ \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1\alpha}^{(2,0)} \left(\frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\alpha}^{(1)}} + \frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\beta}^{(1)}} + \sum_{\gamma \in \mathcal{H}} \left(\frac{p_{\alpha\beta\gamma}^{(3)}}{\Sigma_{\gamma}^{(1)}} - \frac{p_{\alpha\gamma}^{(2)}p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)^2}} \right) \Sigma_{\gamma}^{(1)} \right) \Sigma_{\beta\alpha_2\alpha_3}^{(3)} \\
&+ \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1\alpha}^{(2,0)} \sum_{\gamma \in \mathcal{H}} \left[\frac{p_{\alpha\beta\gamma}^{(3)}}{\Sigma_{\alpha}^{(1)}} + \frac{p_{\alpha\beta\gamma}^{(3)}}{\Sigma_{\beta}^{(1)}} + \frac{p_{\alpha\beta\gamma}^{(3)}}{\Sigma_{\gamma}^{(1)}} - \frac{p_{\beta\alpha}^{(2)}p_{\alpha\gamma}^{(2)}}{\Sigma_{\alpha}^{(1)^2}} - \frac{p_{\alpha\beta}^{(2)}p_{\beta\gamma}^{(2)}}{\Sigma_{\beta}^{(1)^2}} - \frac{p_{\alpha\gamma}^{(2)}p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)^2}} \right. \\
&\quad \left. + \sum_{\delta \in \mathcal{H}} \left(\frac{p_{\alpha\beta\gamma\delta}^{(4)}}{\Sigma_{\delta}^{(1)}} - \frac{p_{\alpha\beta\delta}^{(3)}p_{\delta\gamma}^{(2)}}{\Sigma_{\delta}^{(1)^2}} - \frac{p_{\alpha\delta\gamma}^{(3)}p_{\delta\beta}^{(2)}}{\Sigma_{\delta}^{(1)^2}} - \frac{p_{\delta\beta\gamma}^{(3)}p_{\delta\alpha}^{(2)}}{\Sigma_{\delta}^{(1)^2}} \right. \right. \\
&\quad \left. \left. + 2 \frac{p_{\alpha\delta}^{(2)}p_{\beta\delta}^{(2)}p_{\gamma\delta}^{(2)}}{\Sigma_{\delta}^{(1)^3}} \right) \Sigma_{\delta}^{(1)} \right] \Sigma_{\beta\alpha_2}^{(2)} \Sigma_{\gamma\alpha_3}^{(2)} = \Sigma_{\alpha_1\alpha_2\alpha_3}^{(3)}. \tag{100}
\end{aligned}$$

In writing this expression we have used the results (83), (85), (86), (87) and (89) of the previous sections as well as the fact that, since $\mathbf{p}^{(1)} = \Sigma^{(1)}$, for any $\alpha, \beta, \gamma, \delta \in \mathcal{H}$ it is

$$p_{\alpha\beta\gamma\delta}'''(\Sigma^{(1)}) = p_{\alpha\beta\gamma\delta}^{(4)}. \tag{101}$$

We have also defined

$$\Sigma_{\alpha\beta\gamma}^{(3,0)} = \Sigma_{\alpha\gamma}^{(2)}\delta_{\alpha\beta} - \left(\Sigma_{\alpha\gamma}^{(2)}\Sigma_{\beta}^{(1)} + \Sigma_{\alpha}^{(1)}\Sigma_{\beta\gamma}^{(2)} \right) \chi_{\alpha\beta}, \tag{102}$$

$\Sigma^{(3,0)}$ in a compact notation, which is, formally, the third cumulant of an uncorrelated multinomial probability density with the first two cumulants equal to $\Sigma^{(1)}$ and $\Sigma^{(2)}$, respectively. According to equation (88), in the first two lines of (100) we have $\sum_{\gamma \in \mathcal{H}} p_{\alpha\beta\gamma}^{(3)} = 0$ and in the fourth line $\sum_{\delta \in \mathcal{H}} p_{\alpha\beta\gamma\delta}^{(4)} = 0$, so that equation (100) becomes

$$\begin{aligned}
&\sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \sum_{\gamma \in \mathcal{H}} \left(\Lambda_{\alpha_1\alpha}^{(2,0)}\Sigma_{\beta\alpha_2}^{(2)}\Sigma_{\gamma\alpha_3}^{(2)} + \Sigma_{\alpha_1\alpha}^{(2,0)}\Lambda_{\beta\alpha_2}^{(2)}\Sigma_{\gamma\alpha_3}^{(2)} + \Sigma_{\alpha_1\alpha}^{(2,0)}\Sigma_{\beta\alpha_2}^{(2)}\Lambda_{\gamma\alpha_3}^{(2)} \right) p_{\alpha\beta\gamma}^{(3)} \\
&= \Delta_{\alpha_1\alpha_2\alpha_3}, \tag{103}
\end{aligned}$$

where we have introduced the matrices $\Lambda^{(2)}$ and $\Lambda^{(2,0)}$ with components $\alpha, \beta \in \mathcal{H}$

$$\Lambda_{\alpha\beta}^{(2)} = \frac{1}{\Sigma_{\alpha}^{(1)}} \left(\Sigma_{\alpha\beta}^{(2)} - \sum_{\gamma \in \mathcal{H}} p_{\alpha\gamma}^{(2)}\Sigma_{\gamma\beta}^{(2)} \right), \tag{104}$$

$$\Lambda_{\alpha\beta}^{(2,0)} = \left(\Sigma_{\alpha\beta}^{(2,0)} - \sum_{\gamma \in \mathcal{H}} \Sigma_{\alpha\gamma}^{(2,0)}p_{\gamma\beta}^{(2)} \right) \frac{1}{\Sigma_{\beta}^{(1)}}, \tag{105}$$

and the tensor Δ with components $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{H}$

$$\Delta_{\alpha_1\alpha_2\alpha_3} = \Sigma_{\alpha_1\alpha_2\alpha_3}^{(3)} - \Sigma_{\alpha_1\alpha_2\alpha_3}^{(3,0)}$$

$$\begin{aligned}
& - \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1 \alpha \alpha_3}^{(3,0)} \left(\frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\alpha}^{(1)}} + \frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\beta}^{(1)}} - \sum_{\gamma \in \mathcal{H}} \frac{p_{\alpha\gamma}^{(2)} p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)}} \right) \Sigma_{\beta\alpha_2}^{(2)} \\
& - \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1 \alpha}^{(2,0)} \left(\frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\alpha}^{(1)}} + \frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\beta}^{(1)}} - \sum_{\gamma \in \mathcal{H}} \frac{p_{\alpha\gamma}^{(2)} p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)}} \right) \Sigma_{\beta\alpha_2 \alpha_3}^{(3)} \\
& + \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \sum_{\gamma \in \mathcal{H}} \Sigma_{\alpha_1 \alpha}^{(2,0)} \left(\frac{p_{\alpha\beta}^{(2)} p_{\alpha\gamma}^{(2)}}{\Sigma_{\alpha}^{(1)^2}} + \frac{p_{\alpha\beta}^{(2)} p_{\beta\gamma}^{(2)}}{\Sigma_{\beta}^{(1)^2}} + \frac{p_{\alpha\gamma}^{(2)} p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)^2}} \right. \\
& \quad \left. - 2 \sum_{\delta \in \mathcal{H}} \frac{p_{\alpha\delta}^{(2)} p_{\beta\delta}^{(2)} p_{\gamma\delta}^{(2)}}{\Sigma_{\delta}^{(1)^2}} \right) \Sigma_{\beta\alpha_2}^{(2)} \Sigma_{\gamma\alpha_3}^{(2)}. \tag{106}
\end{aligned}$$

Note that, once $\mathbf{p}^{(1)}$ and $\mathbf{p}^{(2)}$ have been determined, this tensor can be considered known.

Equation (103) is, for $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{H}$, an overdetermined linear system of m^3 equations in the unknown $\mathbf{p}^{(3)}$ which, according to the sum rules (88), has \hat{m}^3 independent components. To find $\mathbf{p}^{(3)}$ we proceed as in the previous section. Let $\hat{\mathbf{p}}^{(3)}$ be the tensor with components $\hat{p}_{\alpha\beta\gamma}^{(3)} = p_{\alpha\beta\gamma}^{(3)}$, $\alpha, \beta, \gamma \in \hat{\mathcal{H}}$, and consider the \hat{m}^3 equations (103) with $\alpha_1, \alpha_2, \alpha_3 \in \hat{\mathcal{H}}$. Observing that $\sum_{\alpha \in \mathcal{H}} = \sum_{\alpha \in \hat{\mathcal{H}}} + \sum_{\alpha \in \{V_*, T_*, \lambda_*\}}$ and using the sum rules (88), we rewrite equation (103) as

$$\begin{aligned}
& \sum_{\alpha \in \hat{\mathcal{H}}} \sum_{\beta \in \hat{\mathcal{H}}} \sum_{\gamma \in \hat{\mathcal{H}}} \left(\tilde{\Lambda}_{\alpha_1 \alpha}^{(2,0)} \tilde{\Sigma}_{\beta\alpha_2}^{(2)} \tilde{\Sigma}_{\gamma\alpha_3}^{(2)} + \tilde{\Sigma}_{\alpha_1 \alpha}^{(2,0)} \tilde{\Lambda}_{\beta\alpha_2}^{(2)} \tilde{\Sigma}_{\gamma\alpha_3}^{(2)} + \tilde{\Sigma}_{\alpha_1 \alpha}^{(2,0)} \tilde{\Sigma}_{\beta\alpha_2}^{(2)} \tilde{\Lambda}_{\gamma\alpha_3}^{(2)} \right) \hat{p}_{\alpha\beta\gamma}^{(3)} \\
& = \Delta_{\alpha_1 \alpha_2 \alpha_3}, \tag{107}
\end{aligned}$$

where we have introduced the matrices $\tilde{\Lambda}^{(2)}$ and $\tilde{\Sigma}^{(2,0)}$ with components $\alpha, \beta \in \hat{\mathcal{H}}$

$$\tilde{\Lambda}_{\alpha\beta}^{(2)} = \Lambda_{\alpha\beta}^{(2)} - \Lambda_{\alpha_*\beta}^{(2)}, \tag{108}$$

$$\tilde{\Lambda}_{\alpha\beta}^{(2,0)} = \Lambda_{\alpha\beta}^{(2,0)} - \Lambda_{\alpha\beta_*}^{(2,0)}, \tag{109}$$

whereas the matrices $\tilde{\Sigma}^{(2)}$ and $\tilde{\Sigma}^{(2,0)}$ are defined by (94) and (96).

The system of equations (107) with $\alpha_1, \alpha_2, \alpha_3 \in \hat{\mathcal{H}}$ is a linear tensorial equation in the unknown $\hat{\mathbf{p}}^{(3)}$. It can be solved by vectorization, i.e. by introducing a bijective map between the set $\hat{\mathcal{H}}^3$ and the integers $\{1, 2, \dots, \hat{m}^3\}$. Let $n(\alpha) : \hat{\mathcal{H}} \mapsto \{1, 2, \dots, \hat{m}\}$ be some ordering of the elements $\alpha \in \hat{\mathcal{H}}$ and $n^{-1} : \{1, 2, \dots, \hat{m}\} \mapsto \hat{\mathcal{H}}$ its inverse. We define the integer map $i(\alpha, \beta, \gamma) : \hat{\mathcal{H}}^3 \mapsto \{1, 2, \dots, \hat{m}^3\}$ by

$$i(\alpha, \beta, \gamma) = (n(\alpha) - 1)\hat{m}^2 + (n(\beta) - 1)\hat{m} + n(\gamma), \tag{110}$$

so that the inverse triplet $(\alpha(i), \beta(i), \gamma(i))$ is given by

$$\alpha(i) = n^{-1} \left(\left\lfloor \frac{i - 0^+}{\hat{m}^2} \right\rfloor + 1 \right), \tag{111}$$

$$\beta(i) = n^{-1} \left(\left\lfloor \frac{i - (n(\alpha(i)) - 1)\hat{m}^2 - 0^+}{\hat{m}} \right\rfloor + 1 \right), \tag{112}$$

$$\gamma(i) = n^{-1} \left(i - (n(\alpha(i)) - 1)\hat{m}^2 - (n(\beta(i)) - 1)\hat{m} \right), \tag{113}$$

where 0^+ is a positive infinitesimal. Instead of equation (107) with $\alpha_1, \alpha_2, \alpha_3 \in \hat{\mathcal{H}}$, we thus obtain the equivalent system

$$\sum_{j=1}^{\hat{m}^3} Q_{ij} \hat{p}_j^{(3)} = \Delta_i, \quad i = 1, 2, \dots, \hat{m}^3, \quad (114)$$

where we have defined

$$Q_{ij} = \tilde{\Lambda}_{\alpha(i)\alpha(j)}^{(2,0)} \tilde{\Sigma}_{\beta(j)\beta(i)}^{(2)} \tilde{\Sigma}_{\gamma(j)\gamma(i)}^{(2)} + \tilde{\Sigma}_{\alpha(i)\alpha(j)}^{(2,0)} \tilde{\Lambda}_{\beta(j)\beta(i)}^{(2)} \tilde{\Sigma}_{\gamma(j)\gamma(i)}^{(2)} + \tilde{\Sigma}_{\alpha(i)\alpha(j)}^{(2,0)} \tilde{\Sigma}_{\beta(j)\beta(i)}^{(2)} \tilde{\Lambda}_{\gamma(j)\gamma(i)}^{(2)}, \quad (115)$$

as well as $\hat{p}_j^{(3)} = \hat{p}_{\alpha(j)\beta(j)\gamma(j)}^{(3)}$ and $\Delta_i = \Delta_{\alpha(i)\beta(i)\gamma(i)}$. Equation (114) is a linear matrix equation which can be solved by standard methods, e.g. LU-factorization [19].

Once $\hat{\mathbf{p}}^{(3)}$ has been found, the complete set of third order perturbative parameters $\mathbf{p}^{(3)}$ is obtained using the sum rules (88) for $k = 3$.

3.6. Some considerations on higher orders

In the previous subsections we have evaluated the perturbative parameters $\mathbf{p}^{(k)}$ for the first three perturbative orders $k = 1, 2, 3$. In this section we will show that for any $k \geq 3$ the equations which determine $\mathbf{p}^{(k)}$ are *i*) linear tensorial equations *ii*) depending only on the parameters $\mathbf{p}^{(j)}$ with $j \leq k$. This ensures that the perturbative parameters are, in principle, solvable at all orders and that their value is independent of the integer $k_{\max} - 1$ at which we decide to truncate the series (49). The evaluation of $\mathbf{p}^{(4)}$ is outlined in Appendix B.

To demonstrate the properties *i*) and *ii*), first of all let us point up why they hold in the analyzed case $k = 3$. The tensor $p_{\alpha_1\alpha_2\alpha_3}^{(3)}$ is determined by the system of equations (55) the structure of which is established, see equation (73), by the third order derivatives $\partial^3 \phi_{\text{sp}}(\mathbf{J}) / \partial J_{\alpha_1} \partial J_{\alpha_2} \partial J_{\alpha_3}$ evaluated at $\mathbf{J} = \mathbf{0}$. According to equation (80), these derivatives contain rational combinations of the functions $\mathbf{p}(\boldsymbol{\nu})$ and their derivatives $\mathbf{p}'(\boldsymbol{\nu})$, $\mathbf{p}''(\boldsymbol{\nu})$ and $\mathbf{p}'''(\boldsymbol{\nu})$, a compact notation to indicate respectively the components (49), (67), (79) and (82), evaluated at $\boldsymbol{\nu} = \boldsymbol{\nu}^{\text{sp}}$. When we set $\mathbf{J} = \mathbf{0}$, since $\boldsymbol{\nu}^{\text{sp}}|_{\mathbf{J}=\mathbf{0}} = \boldsymbol{\Sigma}^{(1)}$ and $\mathbf{p}^{(1)} = \boldsymbol{\Sigma}^{(1)}$, we have

$$\mathbf{p}(\boldsymbol{\nu}^{\text{sp}})|_{\mathbf{J}=\mathbf{0}} = \mathbf{p}^{(1)}, \quad (116)$$

$$\mathbf{p}'(\boldsymbol{\nu}^{\text{sp}})|_{\mathbf{J}=\mathbf{0}} = \mathbf{p}^{(2)}, \quad (117)$$

$$\mathbf{p}''(\boldsymbol{\nu}^{\text{sp}})|_{\mathbf{J}=\mathbf{0}} = \mathbf{p}^{(3)}, \quad (118)$$

$$\mathbf{p}'''(\boldsymbol{\nu}^{\text{sp}})|_{\mathbf{J}=\mathbf{0}} = \mathbf{p}^{(4)}, \quad (119)$$

and so on. Notice that in the third order derivatives (80) there is only one term which contains \mathbf{p}''' , namely

$$\sum_{\alpha, \beta, \gamma, \delta \in \mathcal{H}} (\nu_{\alpha_1}^{\text{sp}} \delta_{\alpha_1\alpha} - \nu_{\alpha_1}^{\text{sp}} \nu_{\alpha}^{\text{sp}} \chi_{\alpha_1\alpha}) \frac{p_{\alpha\beta\gamma\delta}'''(\boldsymbol{\nu}^{\text{sp}})}{p_{\delta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\delta}^{\text{sp}} \frac{\partial \nu_{\beta}^{\text{sp}}}{\partial J_{\alpha_2}} \frac{\partial \nu_{\gamma}^{\text{sp}}}{\partial J_{\alpha_3}}. \quad (120)$$

When this term is evaluated at $\mathbf{J} = \mathbf{0}$, the factors $\nu_{\delta}^{\text{sp}}|_{\mathbf{J}=\mathbf{0}}$ and $p_{\delta}(\boldsymbol{\nu}^{\text{sp}})|_{\mathbf{J}=\mathbf{0}}$ cancels each other out so that, according to the sum rules (88), we have

$$\sum_{\delta \in \mathcal{H}} p_{\alpha\beta\gamma\delta}'''(\boldsymbol{\nu}^{\text{sp}})|_{\mathbf{J}=\mathbf{0}} = \sum_{\delta \in \mathcal{H}} p_{\alpha\beta\gamma\delta}^{(4)} = 0. \quad (121)$$

As a consequence, the system of equations which determines $\mathbf{p}^{(3)}$ does not contain $\mathbf{p}^{(4)}$. The property is immediately extended to higher orders. The system of equations which determines $p_{\alpha_1\alpha_2\alpha_3\alpha_4}^{(4)}$ depends on the fourth order derivatives $\partial^4\phi_{\text{sp}}(\mathbf{J})/\partial J_{\alpha_1}\partial J_{\alpha_2}\partial J_{\alpha_3}\partial J_{\alpha_4}$ and these contain \mathbf{p} , \mathbf{p}' , \mathbf{p}'' , \mathbf{p}''' and \mathbf{p}'''' . The fourth order derivative \mathbf{p}'''' may appear only in the term

$$\sum_{\alpha,\beta,\gamma,\delta,\varepsilon\in\mathcal{H}} (\nu_{\alpha_1}^{\text{sp}}\delta_{\alpha_1\alpha} - \nu_{\alpha_1}^{\text{sp}}\nu_{\alpha}^{\text{sp}}\chi_{\alpha_1\alpha}) \frac{p_{\alpha\beta\gamma\delta\varepsilon}''''(\boldsymbol{\nu}^{\text{sp}})}{p_{\delta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\delta}^{\text{sp}} \frac{\partial \nu_{\beta}^{\text{sp}}}{\partial J_{\alpha_2}} \frac{\partial \nu_{\gamma}^{\text{sp}}}{\partial J_{\alpha_3}} \frac{\partial \nu_{\varepsilon}^{\text{sp}}}{\partial J_{\alpha_4}}, \quad (122)$$

obtained differentiating the factor $p_{\alpha\beta\gamma\delta}'''(\boldsymbol{\nu}^{\text{sp}})$ of (120) with respect to J_{α_4} . When evaluated at $\mathbf{J} = \mathbf{0}$, equation (122) vanishes and we find that no $\mathbf{p}^{(5)}$ terms are contained in the system of equations for $\mathbf{p}^{(4)}$. Iterating, we conclude that the property *ii*) is valid at any higher order.

Now we focus on the property *i*). The system of equations which determines $\mathbf{p}^{(3)}$ is linear in the unknown tensor simply because the third order derivatives $\partial^3\phi_{\text{sp}}(\mathbf{J})/\partial J_{\alpha_1}\partial J_{\alpha_2}\partial J_{\alpha_3}$ are linear in \mathbf{p}'' , see equation (80). By further differentiating (80) with respect to J_{α_4} there is no possibility to generate a term nonlinear in \mathbf{p}''' , e.g. quadratic. In fact, this would amount to have in $\partial^3\phi_{\text{sp}}(\mathbf{J})/\partial J_{\alpha_1}\partial J_{\alpha_2}\partial J_{\alpha_3}$ a term containing the product or the ratio of the components of \mathbf{p}''' and \mathbf{p}'' . However, we have seen that the only term of (80) containing \mathbf{p}''' is given by (120). We conclude that the system of equations which determines $\mathbf{p}^{(4)}$ is linear and, iterating, the same holds at any higher order.

4. Evaluation of the ground-state energy

We have already seen that the ground state energy E_0 is the unique solution of the scalar equation

$$\lim_{N\rightarrow\infty} \frac{1}{N} \log I_N(E_0) = 0, \quad E_0 \leq V_{\min}, \quad (123)$$

where

$$I_N(E_0) = \int d(N\boldsymbol{\nu}) \mathcal{P}_N(N\boldsymbol{\nu}) e^{(N\boldsymbol{\nu}, \mathbf{u}(E_0))}, \quad (124)$$

and

$$\mathbf{u}^T(E_0) = (\dots - \log(-E_0 + V) \dots; \dots \log T \dots; \dots \log \lambda \dots). \quad (125)$$

In this section we want to determine E_0 when the probability density $\mathcal{P}_N(N\boldsymbol{\nu})$ is given by the multinomial perturbative scheme (48-49).

For N large we have already calculated the integral (124). In fact, $I_N(E_0)$ coincides with the generating function $Z_N(\mathbf{J})$ studied in section 3.1 provided we choose the source $\mathbf{J} = \mathbf{u}(E_0)$. Therefore, equation (123) is equivalent to

$$\phi_{\text{sp}}(\mathbf{u}(E_0)) = 0, \quad E_0 \leq V_{\min}, \quad (126)$$

where $\phi_{\text{sp}}(\mathbf{u}(E_0))$ is given by (72) with $\mathbf{J} = \mathbf{u}(E_0)$. Of course, also the saddle-point frequencies (71) which appear in (72) must be evaluated with the same choice of \mathbf{J} . We

conclude that the ground-state energy E_0 is obtained, together with the saddle-point frequencies $\boldsymbol{\nu}^{\text{sp}}$, as the solution of the following system of nonlinear coupled equations

$$\sum_{V \in \mathcal{V}} \frac{\tilde{p}_V(\boldsymbol{\nu}^{\text{sp}})}{-E_0 + V} = \frac{e^{\sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{p_{\beta}(\boldsymbol{\nu}^{\text{sp}})} \nu_{\alpha}^{\text{sp}} \nu_{\beta}^{\text{sp}}}}{(\sum_{T \in \mathcal{T}} \tilde{p}_T(\boldsymbol{\nu}^{\text{sp}}) T) (\sum_{\lambda \in \mathcal{L}} \tilde{p}_{\lambda}(\boldsymbol{\nu}^{\text{sp}}) \lambda)}, \quad E_0 \leq V_{\min}, \quad (127)$$

$$\nu_{\alpha}^{\text{sp}} = \frac{\tilde{p}_{\alpha}(\boldsymbol{\nu}^{\text{sp}}) e^{u_{\alpha}(E_0)}}{\sum_{\alpha' \in \mathcal{H}_{\alpha}} \tilde{p}_{\alpha'}(\boldsymbol{\nu}^{\text{sp}}) e^{u_{\alpha'}(E_0)}}, \quad \alpha \in \mathcal{H}, \quad (128)$$

where we have defined

$$\tilde{p}_{\alpha}(\boldsymbol{\nu}) = p_{\alpha}(\boldsymbol{\nu}) e^{\sum_{\beta \in \mathcal{H}} \frac{p'_{\alpha\beta}(\boldsymbol{\nu})}{p_{\beta}(\boldsymbol{\nu})} \nu_{\beta}}, \quad (129)$$

and we recall that

$$p_{\alpha}(\boldsymbol{\nu}) = \sum_{n=0}^{k_{\max}-1} \frac{1}{n!} \sum_{\beta_1 \in \mathcal{H}} \cdots \sum_{\beta_n \in \mathcal{H}} p_{\alpha\beta_1 \dots \beta_n}^{(n+1)} \left(\nu_{\beta_1} - \Sigma_{\beta_1}^{(1)} \right) \cdots \left(\nu_{\beta_n} - \Sigma_{\beta_n}^{(1)} \right),$$

$$p'_{\alpha\beta}(\boldsymbol{\nu}) = \sum_{n=1}^{k_{\max}-1} \frac{1}{(n-1)!} \sum_{\gamma_1 \in \mathcal{H}} \cdots \sum_{\gamma_{n-1} \in \mathcal{H}} p_{\alpha\beta\gamma_1 \dots \gamma_{n-1}}^{(n+1)} \times \left(\nu_{\gamma_1} - \Sigma_{\gamma_1}^{(1)} \right) \cdots \left(\nu_{\gamma_{n-1}} - \Sigma_{\gamma_{n-1}}^{(1)} \right).$$

Note that the above system of equations is valid at any perturbative order. More precisely, for different choices of k_{\max} only the functions $p_{\alpha}(\boldsymbol{\nu})$ and $p'_{\alpha\beta}(\boldsymbol{\nu})$ are to be modified whereas the structure of the equations (127-128) remain unchanged.

At the lowest perturbative order $k_{\max} = 1$, we have $\mathbf{p}(\boldsymbol{\nu}) = \boldsymbol{\Sigma}^{(1)}$, $\mathbf{p}'(\boldsymbol{\nu}) = \mathbf{0}$ and $\tilde{\mathbf{p}}(\boldsymbol{\nu}) = \boldsymbol{\Sigma}^{(1)}$. It follows that equations (127) and (128) can be solved separately. Equation (127) reduces to (44), i.e. E_0 is the solution of

$$\sum_{V \in \mathcal{V}} \frac{\Sigma_V^{(1)}}{-E_0 + V} = \frac{1}{\left(\sum_{T \in \mathcal{T}} \Sigma_T^{(1)} T \right) \left(\sum_{\lambda \in \mathcal{L}} \Sigma_{\lambda}^{(1)} \lambda \right)}, \quad E_0 \leq V_{\min}. \quad (130)$$

This equation can be straightforwardly solved by bisection method. Once E_0 is found, the saddle-point frequencies are given by

$$\nu_V^{\text{sp}} = \frac{\Sigma_V^{(1)} (-E_0 + V)^{-1}}{\sum_{V' \in \mathcal{V}} \Sigma_{V'}^{(1)} (-E_0 + V')^{-1}}, \quad V \in \mathcal{V}, \quad (131)$$

$$\nu_T^{\text{sp}} = \frac{\Sigma_T^{(1)} T}{\sum_{T' \in \mathcal{T}} \Sigma_{T'}^{(1)} T'}, \quad T \in \mathcal{T}, \quad (132)$$

$$\nu_{\lambda}^{\text{sp}} = \frac{\Sigma_{\lambda}^{(1)} \lambda}{\sum_{\lambda' \in \mathcal{L}} \Sigma_{\lambda'}^{(1)} \lambda'}, \quad \lambda \in \mathcal{L}. \quad (133)$$

At higher perturbative orders $k_{\max} > 1$, once the perturbative parameters $\mathbf{p}^{(k)}$ for $k = 1, \dots, k_{\max}$ have been determined from the corresponding exact cumulants $\boldsymbol{\Sigma}^{(1)}, \dots, \boldsymbol{\Sigma}^{(k_{\max})}$, the system of equations (127) and (128) can be solved numerically by a globally convergent multidimensional Newton–Raphson method [19]. Of course, we

have to conjecture some initial value of (E_0, ν^{sp}) which is not too far from the solution. Actually, this may not represent a problem in the spirit of our perturbative approach according to which the uncorrelated multinomial probability density can roughly capture the features of the system considered. Therefore, we propose to use the solution of (130) and the values (131-133) as an initial guess for (E_0, ν^{sp}) .

5. Numerical results for the Hubbard model

In this section we apply the multinomial perturbative scheme to study some example systems. We will focus on the Hubbard model with spin $\frac{1}{2}$ hard-core bosons. The case of fermions will be considered in a paper dedicated to the sign problem.

The Hubbard model is one of the simplest models displaying the real word features of a many-body system. It plays essentially the same role in the problem of the electron correlations as the Ising model in the problem of spin correlations. The model describes interacting particles, bosons or fermions, in a lattice, and the corresponding Hamiltonian operator consists of two terms: a kinetic term allowing for hopping of particles among the sites of the lattice and a potential term consisting of an on-site interaction. The model was originally proposed by Hubbard [20] to describe electrons in solids and has since been the focus of particular interest as a model for high-temperature superconductivity. Recently a large interest has been devoted also to the properties of the 2D Hubbard model on the honeycomb lattice, as a basic model for the description of graphene [21].

Let us consider the first-neighbour uniform (FNU) Hubbard model defined by the Hamiltonian

$$\hat{H} = -\eta \sum_{(i,j) \in \Gamma} \sum_{\sigma=\uparrow\downarrow} \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} + \hat{c}_{j\sigma}^\dagger \hat{c}_{i\sigma} \right) + \gamma \sum_{i \in \Lambda} \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\uparrow} \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow}, \quad (134)$$

where $\Lambda \subset \mathbb{Z}^d$ is a finite d -dimensional lattice with $|\Lambda|$ ordered sites, $\Gamma = \{(i, j) : i < j \in \Lambda \text{ and } i, j \text{ are first neighbour}\}$, and $\hat{c}_{i\sigma}$ a commuting destruction operator at site $i \in \Lambda$ and spin index $\sigma = \uparrow\downarrow$ with the property $\hat{c}_{i\sigma}^2 = 0$ (hard-core boson destruction operator). Hopping strengths are uniform through the lattice and described by the parameter $\eta > 0$. Also the on-site interactions are independent of the site index and their value is fixed by the parameter $\gamma \geq 0$. The system is described in terms of Fock states labelled by the configurations $\mathbf{n} = (n_{1\uparrow}, n_{1\downarrow}, \dots, n_{|\Lambda|\uparrow}, n_{|\Lambda|\downarrow})$ with $n_{i\sigma} = 0, 1$. In this \mathbf{n} -representation, the on-site Hubbard interaction is the diagonal potential operator \hat{V} with matrix elements

$$V_{\mathbf{n}, \mathbf{n}} = V_{\mathbf{n}} = \gamma \sum_{i \in \Lambda} n_{i\uparrow} n_{i\downarrow}, \quad (135)$$

whereas the matrix elements of the hopping operator are

$$K_{\mathbf{n}, \mathbf{n}'} = -\eta \sum_{(i,j) \in \Gamma} \sum_{\sigma=\uparrow\downarrow} \delta_{\mathbf{n}, \mathbf{n}' \oplus \mathbf{1}_{i\sigma} \oplus \mathbf{1}_{j\sigma}}, \quad (136)$$

where $\mathbf{1}_{i\sigma} = (0, \dots, 0, 1_{i\sigma}, 0, \dots, 0)$ and \oplus means mod 2 addition. By comparing (136) with (2) and observing that $\sum_{(i,j) \in \Gamma} \sum_{\sigma=\uparrow\downarrow} \delta_{\mathbf{n}, \mathbf{n}' \oplus \mathbf{1}_{i\sigma} \oplus \mathbf{1}_{j\sigma}} = 0, 1$, the unit value being

obtained only if $\mathbf{n}' \in A(\mathbf{n})$, where

$$A(\mathbf{n}) = \{\mathbf{n} \oplus \mathbf{1}_{i\sigma} \oplus \mathbf{1}_{j\sigma} : (i, j) \in \Gamma, \sigma = \uparrow\downarrow \text{ with } n_{i\sigma} \oplus n_{j\sigma} = 1\} \quad (137)$$

is the set of the configurations connected to \mathbf{n} by the hopping of one particle, we have

$$\eta_{\mathbf{n}, \mathbf{n}'} = \eta, \quad \lambda_{\mathbf{n}, \mathbf{n}'} = \begin{cases} 1, & \mathbf{n}' \in A(\mathbf{n}), \\ 0, & \text{otherwise.} \end{cases} \quad (138)$$

In the following we will consider two dimensional square lattices having $L_x \times L_y$ sites and containing N_p particles per spin. Periodic boundary conditions will be imposed. At the densities $N_p/(L_x L_y) \leq 1/2$ took into consideration, the set of all possible different values of the potential variables (7) during an infinitely long random walk corresponds to the set of all possible different values of $V_{\mathbf{n}}$, the matrix elements (135), over the configuration space. It is straightforward to see that $\mathcal{V} = \{0, \gamma, 2\gamma, \dots, N_p \gamma\}$. The set of all possible different values of the hopping variables (17) during an infinitely long random walk, corresponds to the set of all possible different values of $T_{\mathbf{n}} = \eta |A(\mathbf{n})|$ over the configuration space, with $A(\mathbf{n})$ given by (137). The determination of \mathcal{T} depends both on the number of particles and on the lattice size. For instance, in a lattice 2×3 we have $\mathcal{T} = \{12\eta, 16\eta, 20\eta\}$ with $N_p = 3$ and $\mathcal{T} = \{8\eta, 10\eta, 12\eta, 14\eta, 16\eta\}$ with $N_p = 2$. For hard-core bosons the set of all possible phase variables is $\mathcal{L} = \{1\}$.

Once we have determined the sets \mathcal{V} , \mathcal{T} and \mathcal{L} , the asymptotic rescaled cumulants $\Sigma^{(k)}$ associated to considered Hubbard system can be measured as explained in [11]. Note that these cumulants are unaffected by a change of the parameters η and γ of the Hamiltonian (134), whereas the label sets \mathcal{V} , \mathcal{T} and \mathcal{L} keep their cardinalities under the same change. These data are input into equations (127-128) to find the ground-state energy E_0 . The results obtained by using cumulants up to order $k_{\max} = 1, 2, 3, 4$ are shown in figure 2 as a function of the ratio γ/η in the case of a lattice with 2×3 sites and $N_p = 2, 3$ particles per spin. In the same figure we also depict the values of E_0 determined by exact numerical diagonalization. The curves obtained for $k_{\max} = 1$ coincide with the uncorrelated multinomial prediction and, as already noted, their behavior is only qualitatively correct. Already at $k_{\max} = 2$, the quantitative agreement with the exact ground-state energy becomes impressive at least for values of γ/η not too large. By further increasing k_{\max} , the quantitative agreement gradually improves in the whole range of γ/η which, note the horizontal log scale, goes from the limit $\|\hat{V}\| \ll \|\hat{K}\|$ to the opposite one $\|\hat{V}\| \gg \|\hat{K}\|$. In the case with $N_p = 2$ particles per spin, the curve $E_0(\gamma)$ obtained with $k_{\max} = 4$ is indistinguishable from the reported exact values. In the case with $N_p = 3$ particles per spin, the nonlinear equations (127-128) do not admit a solution with $k_{\max} = 3, 4$ when γ/η is larger than a threshold value, namely $\gamma/\eta \simeq 31.6$ for $k_{\max} = 3$ and $\gamma/\eta \simeq 2.5$ for $k_{\max} = 4$. More precisely, above this threshold equations (127-128) admit only solutions with real negative or complex frequencies ν . As discussed at the beginning of section 3, these are meaningless results which make the perturbative scheme invalid at the order considered.

In figure 3 we show the results obtained with a system of larger size, namely a lattice 4×4 with $N_p = 2, 4, 5, 8$ particles. In this case, the number of configurations is




Figure 2. Ground-state energy per particle for the 2×3 FNU hard-core boson Hubbard model with periodic boundary conditions versus the interaction strength γ with $N_p = 2$ and $N_p = 3$ particles per spin. We compare the results from exact numerical diagonalization (\times) with those from present multinomial perturbative scheme by using cumulants up to order $k_{\max} = 1, 2, 3, 4$ (dotted, dashed, dot-dashed, solid lines, respectively).

so large, namely

$$M = \left(\frac{(L_x L_y)!}{N_p! (L_x L_y - N_p)!} \right)^2, \quad (139)$$

that an exact numerical diagonalization of the Hamiltonian (134) is unfeasible. Therefore, we have compared the values of E_0 predicted by the present multinomial perturbative scheme with those obtained by a Monte Carlo simulation [7]. The conclusions that we reach from the analysis of figure 3 are similar to those we noticed after figure 2. However, for this larger system we see that the parity of the order k_{\max} may influence the quality of the approximation, a fact which is not surprising. For instance, in the case with $N_p = 5$ it is evident that the curve $E_0(\gamma)$ obtained with $k_{\max} = 3$ is not better than that obtained with $k_{\max} = 2$. However, as evidenced in the enlargement shown in figure 4, the results obtained with $k_{\max} = 4$ are more accurate than those with $k_{\max} = 2$, at least in the γ/η range where equations (127-128) admit the $k_{\max} = 4$ solution.

6. Conclusions

In the framework of the probabilistic approach previously developed by us to study the ground-state properties of a many-body quantum system, we have introduced a multinomial perturbative scheme which has the following characteristics. At any




Figure 3. As in figure 1 in the case of a 4×4 FNU hard-core boson Hubbard model with periodic boundary conditions and $N_p = 2, 4, 5, 8$ particles per spin. The data indicated by \circ have been obtained by Monte Carlo simulations [7] (the associated statistical errors increase by increasing γ and are of the order of the symbol size at $\gamma \simeq 1000\eta$).




Figure 4. Enlargement of figure 3, case with $N_p = 5$. The perturbative scheme with $k_{\max} = 4$ admits a solution only for $\gamma \lesssim 3.2\eta$ but in this range provides results more accurate than those obtained with $k_{\max} = 1, 2, 3$.

perturbative order, the probability distribution of the potential, hopping and phase multiplicities, whose knowledge would allow for an exact solution of the problem,

is approximated by a multinomial-like distribution with infinitely many statistical moments. By increasing the perturbative order, an increasing number of cumulants of the distribution is made to coincide with the corresponding exact cumulants of the system.

We have tested the proposed perturbative scheme in the case of two-dimensional lattice Hubbard models with spin $\frac{1}{2}$ hard-core bosons. Already at second perturbative order, we find a ground-state energy in good quantitative agreement with the exact one for any value of the ratio γ/η , γ and η being the strengths of the interaction and hopping terms of the Hamiltonian of the system. The agreement improves at higher perturbative orders. At orders ≥ 3 , however, the scheme may not be always consistent, i.e. in some systems a solution for the ground-state energy is found only for values of γ/η smaller than a threshold. As a matter of fact, we observe that in all our test cases at, or near, the $\frac{1}{4}$ particle filling, which is a case of remarkable physical interest, the solution of the perturbative method turns out to exist for all values of the coupling parameter up to the largest explored perturbative order, $k_{\max} = 4$, where it provides stunning results.

The perturbative probability distribution at order k is built up from the knowledge of the first k connected statistical moments of the potential, hopping and phase multiplicities of the system. These cumulants are easily measured by Monte Carlo simulations and are independent of the parameters γ and η of the Hamiltonian. Once the probability distribution is determined at the chosen approximation, the ground-state energy E_0 , or any other correlation function in the ground state of the system, can be found by solving numerically a small system of nonlinear equations. This last job has a computational cost negligible with respect to the determination of the cumulants, which, in turn, has a cost equivalent to a direct Monte Carlo evaluation of E_0 . Thus, the advantage of our approach in comparison to a direct Monte Carlo simulation is remarkable. Different Monte Carlo runs are needed to evaluate E_0 for different values of γ and/or η , whereas in our approach we have to solve each time a small system of nonlinear equations and, *una tantum*, calculate the cumulants.

The present perturbative probabilistic approach is particularly promising for systems affected by the so called sign problem, for which unbiased Monte Carlo simulations of E_0 are impractical. In fact, the statistical evaluation of the cumulants of the potential, hopping and phase multiplicities is unaffected by the sign problem, which remains confined in the expression of the probability distribution and can be tackled by analytical techniques. We plan to discuss the case of fermions in a future paper.

Appendix A. Solution of nonsymmetric algebraic Riccati equations

In section 3.4 we have seen that the parameters $\mathbf{p}^{(2)}$, more precisely the associated reduced matrix $\hat{\mathbf{p}}^{(2)}$, are determined by the NARE (99). In general, the NAREs are defined as the quadratic matrix equations of the kind

$$\mathbf{X}\mathbf{C}\mathbf{X} - \mathbf{A}\mathbf{X} - \mathbf{X}\mathbf{D} + \mathbf{B} = \mathbf{0}, \quad (\text{A.1})$$

where we assume that the unknown \mathbf{X} , as well as the coefficients \mathbf{A} , \mathbf{B} , \mathbf{C} and \mathbf{D} are quadratic matrices of finite size. In this section we illustrate two numerical methods developed to solve equation (A.1). The first one is an iterative method based on a fixed-point technique [22], whereas the second one is a direct method based on the Schür decomposition [18].

Equations (A.1) play an important role in the study of the stochastic fluid models and have been extensively studied. In general, a NARE admits more than one solution. In most stochastic fluid models, the coefficients \mathbf{A} , \mathbf{B} , \mathbf{C} and \mathbf{D} form a super matrix

$$\mathbf{H} = \begin{pmatrix} \mathbf{D} & -\mathbf{C} \\ -\mathbf{B} & \mathbf{A} \end{pmatrix}. \quad (\text{A.2})$$

with the property to be a so called M -matrix [18]. It can be proved that in this case the Schür method provides the minimal non negative solution of the NARE, which is, there, the solution of physical interest.

In our context, \mathbf{H} is not a M -matrix and it is not clear which solution of the NARE (99) has to be considered. We propose to consider the unique solution given by the Schür method. This solution coincides with that obtained by the iterative method in which \mathbf{X} is chosen at the 0-th iteration as the solution of (A.1) with $\mathbf{C} = \mathbf{0}$. Since in our case \mathbf{X} represents the matrix of the perturbative parameters $\hat{\mathbf{p}}^{(2)}$, which we expect to be small, the above proposed solution seems the most natural one.

Appendix A.1. Iterative method

In [22] a class of fixed-point methods is considered to solve equation (A.1). These fixed-point iterations are based on a suitable splitting of the matrices \mathbf{A} and \mathbf{D} , that is $\mathbf{A} = \mathbf{A}_1 - \mathbf{A}_2$ and $\mathbf{D} = \mathbf{D}_1 - \mathbf{D}_2$, and have the form

$$\mathbf{A}_1 \mathbf{X}_{k+1} + \mathbf{X}_{k+1} \mathbf{D}_1 = \mathbf{X}_k \mathbf{C} \mathbf{X}_k + \mathbf{A}_2 \mathbf{X}_k + \mathbf{X}_k \mathbf{D}_2 + \mathbf{B}, \quad (\text{A.3})$$

with $k = 0, 1, 2, \dots$ and $\mathbf{X}_0 = \mathbf{0}$. In particular, for $\mathbf{A}_1 = \mathbf{A}$ and $\mathbf{D}_1 = \mathbf{D}$ we have

$$\mathbf{A} \mathbf{X}_{k+1} + \mathbf{X}_{k+1} \mathbf{D} = \mathbf{X}_k \mathbf{C} \mathbf{X}_k + \mathbf{B}, \quad \mathbf{X}_0 = \mathbf{0}. \quad (\text{A.4})$$

Note that finding \mathbf{X}_{k+1} in terms of \mathbf{X}_k at k -th iteration implies to solve a Sylvester equation. This can be accomplished by vectorization, namely

$$\text{vec}(\mathbf{A} \mathbf{X}_{k+1}) + \text{vec}(\mathbf{X}_{k+1} \mathbf{D}) = \text{vec}(\mathbf{X}_k \mathbf{C} \mathbf{X}_k + \mathbf{B}). \quad (\text{A.5})$$

Using the properties of the vec operator, in particular

$$\text{vec}(\mathbf{A} \mathbf{X}_{k+1}) = (\mathbf{I} \otimes \mathbf{A}) \text{vec}(\mathbf{X}_{k+1}), \quad (\text{A.6})$$

$$\text{vec}(\mathbf{X}_{k+1} \mathbf{D}) = (\mathbf{D}^T \otimes \mathbf{I}) \text{vec}(\mathbf{X}_{k+1}), \quad (\text{A.7})$$

where \otimes indicates the Kronecker product and T the transpose, equation (A.5) is rewritten as

$$[(\mathbf{I} \otimes \mathbf{A}) + (\mathbf{D}^T \otimes \mathbf{I})] \text{vec}(\mathbf{X}_{k+1}) = \text{vec}(\mathbf{X}_k \mathbf{C} \mathbf{X}_k + \mathbf{B}). \quad (\text{A.8})$$

This is a linear matrix equation which can be solved by standard methods, e.g. LU-factorization [19].

The convergence of the full class of iterative schemes (A.3) to a solution \mathbf{X} of (A.1) is ensured by a theorem [22]. In this class, the iterative scheme (A.4) is the most expensive from a computational point of view, but, on the other hand, it has the highest (linear) convergence speed.

Appendix A.2. Schür method

In the following we discuss a different approach to solve equation (A.1) based on the ordered Schür decomposition. This approach was conceived by Laub [17] for a symmetric algebraic Riccati equation and extended by Guo [23] to the study of NAREs.

Let us rewrite the matrix \mathbf{H} associated to the coefficients of (A.1) as

$$\mathbf{H} = \begin{pmatrix} \mathbf{D} & -\mathbf{C} \\ -\mathbf{B} & \mathbf{A} \end{pmatrix} = \begin{pmatrix} \mathbf{H}_{11} & \mathbf{H}_{12} \\ \mathbf{H}_{21} & \mathbf{H}_{22} \end{pmatrix}. \quad (\text{A.9})$$

Note that \mathbf{H} is real in our case. We look for an orthogonal transformation

$$\mathbf{U} = \begin{pmatrix} \mathbf{U}_{11} & \mathbf{U}_{12} \\ \mathbf{U}_{21} & \mathbf{U}_{22} \end{pmatrix}, \quad (\text{A.10})$$

which leaves \mathbf{H} in a semi-ordered real Schür form,

$$\mathbf{U}^T \mathbf{H} \mathbf{U} = \mathbf{S} = \begin{pmatrix} \mathbf{S}_{11} & \mathbf{S}_{12} \\ \mathbf{0} & \mathbf{S}_{22} \end{pmatrix}, \quad (\text{A.11})$$

in which \mathbf{S}_{11} and \mathbf{S}_{22} contain only blocks, denoted \mathbf{s}_{ij} , $i, j = 1, 2, \dots$, of size 1 or 2. The eigenvalues of the 2×2 diagonal blocks \mathbf{s}_{ii} provide the complex conjugated eigenvalues of \mathbf{H} whereas the 1×1 blocks are the real eigenvalues of \mathbf{H} . The diagonal blocks are semi-ordered in the sense that if \mathbf{s}_{ii} , \mathbf{s}_{jj} and \mathbf{s}_{kk} have eigenvalues with positive, null and negative real parts, respectively, then $i < j < k$. It is possible to show that the matrix \mathbf{U}_{11} is invertible [24] and that

$$\mathbf{X} = \mathbf{U}_{21} \mathbf{U}_{11}^{-1} \quad (\text{A.12})$$

solves (A.1). Note that the semi-ordered decomposition (A.11) is unique and so is the solution (A.12). We used the subroutines of LAPACK library [25] to numerically implement the Schür method.

Appendix B. Equations for the perturbative parameters: fourth order

The perturbative parameters $\mathbf{p}^{(4)}$ are determined by the system of equations

$$\langle\langle \nu_{\alpha_1} \nu_{\alpha_2} \nu_{\alpha_3} \nu_{\alpha_4} \rangle\rangle(\mathbf{p}^{(1)}, \mathbf{p}^{(2)}, \mathbf{p}^{(3)}, \mathbf{p}^{(4)}) = \Sigma_{\alpha_1 \alpha_2 \alpha_3, \alpha_4}^{(4)}, \quad (\text{B.1})$$

with $\alpha_1, \alpha_2, \alpha_3, \alpha_4 \in \mathcal{H}$. By using (73) and taking the derivative of (80) with respect to J_{α_4} , the above system can be cast in the form

$$\sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \sum_{\gamma \in \mathcal{H}} \sum_{\delta \in \mathcal{H}} \left(\Lambda_{\alpha_1 \alpha}^{(2,0)} \Sigma_{\beta \alpha_2}^{(2)} \Sigma_{\gamma \alpha_3}^{(2)} \Sigma_{\delta \alpha_4}^{(2)} + \Sigma_{\alpha_1 \alpha}^{(2,0)} \Lambda_{\beta \alpha_2}^{(2)} \Sigma_{\gamma \alpha_3}^{(2)} \Sigma_{\delta \alpha_4}^{(2)} \right)$$

$$+ \Sigma_{\alpha_1\alpha}^{(2,0)} \Sigma_{\beta\alpha_2}^{(2)} \Lambda_{\gamma\alpha_3}^{(2)} \Sigma_{\delta\alpha_4}^{(2)} + \Sigma_{\alpha_1\alpha}^{(2,0)} \Sigma_{\beta\alpha_2}^{(2)} \Sigma_{\gamma\alpha_3}^{(2)} \Lambda_{\delta\alpha_4}^{(2)} \Big) p_{\alpha\beta\gamma\delta}^{(4)} = \Delta_{\alpha_1\alpha_2\alpha_3\alpha_4}, \quad (\text{B.2})$$

where $\Sigma^{(2)}$ is the asymptotic rescaled cumulant of order 2 and the matrices $\Sigma^{(2,0)}$, $\Lambda^{(2)}$ and $\Lambda^{(2,0)}$ are defined by (90), (104) and (105), respectively. The tensor Δ has components $\alpha_1, \alpha_2, \alpha_3, \alpha_4 \in \mathcal{H}$ given by

$$\begin{aligned} \Delta_{\alpha_1\alpha_2\alpha_3\alpha_4} &= \Sigma_{\alpha_1\alpha_2\alpha_3\alpha_4}^{(4)} - \Sigma_{\alpha_1\alpha_2\alpha_3\alpha_4}^{(4,0)} \\ &\quad - \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \left(\Sigma_{\alpha_1\alpha}^{(2,0)} M_{\alpha\beta}^{(2)} \Sigma_{\beta\alpha_2\alpha_3\alpha_4}^{(4)} + \Sigma_{\alpha_1\alpha\alpha_3\alpha_4}^{(4,0)} M_{\alpha\beta}^{(2)} \Sigma_{\beta\alpha_2}^{(2)} \right) \\ &\quad - \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1\alpha\alpha_3}^{(3,0)} \left(M_{\alpha\beta\alpha_4}^{(3)} \Sigma_{\beta\alpha_2}^{(2)} + M_{\alpha\beta}^{(2)} \Sigma_{\beta\alpha_2\alpha_4}^{(3)} \right) \\ &\quad - \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1\alpha\alpha_4}^{(3,0)} \left(M_{\alpha\beta\alpha_3}^{(3)} \Sigma_{\beta\alpha_2}^{(2)} + M_{\alpha\beta}^{(2)} \Sigma_{\beta\alpha_2\alpha_3}^{(3)} \right) \\ &\quad - \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \Sigma_{\alpha_1\alpha}^{(2,0)} \left(M_{\alpha\beta\alpha_3}^{(3)} \Sigma_{\beta\alpha_2\alpha_4}^{(3)} + M_{\alpha\beta\alpha_4}^{(3)} \Sigma_{\beta\alpha_2\alpha_3}^{(3)} \right) \\ &\quad - \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \sum_{\gamma \in \mathcal{H}} \Sigma_{\alpha_1\alpha}^{(2,0)} F_{\alpha\beta\gamma}^{(3)} \Sigma_{\beta\alpha_2}^{(2)} \Sigma_{\gamma\alpha_3\alpha_4}^{(3)} \\ &\quad + \sum_{\alpha \in \mathcal{H}} \sum_{\beta \in \mathcal{H}} \sum_{\gamma \in \mathcal{H}} \sum_{\delta \in \mathcal{H}} \Sigma_{\alpha_1\alpha}^{(2,0)} F_{\alpha\beta\gamma\delta}^{(4)} \Sigma_{\beta\alpha_2}^{(2)} \Sigma_{\gamma\alpha_3}^{(2)} \Sigma_{\delta\alpha_4}^{(2)}, \end{aligned} \quad (\text{B.3})$$

where

$$\Sigma_{\alpha\beta\gamma\delta}^{(4,0)} = \Sigma_{\beta\gamma\delta}^{(3)} \delta_{\alpha\beta} - \left(\Sigma_{\alpha\gamma\delta}^{(3)} \Sigma_{\beta}^{(1)} + \Sigma_{\beta\gamma\delta}^{(3)} \Sigma_{\alpha}^{(1)} + \Sigma_{\alpha\delta}^{(2)} \Sigma_{\beta\gamma}^{(2)} + \Sigma_{\alpha\gamma}^{(2)} \Sigma_{\beta\delta}^{(2)} \right) \chi_{\alpha\beta}, \quad (\text{B.4})$$

$$M_{\alpha\beta}^{(2)} = M_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})|_{\mathbf{J}=\mathbf{0}} = \frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\alpha}^{(1)}} + \frac{p_{\alpha\beta}^{(2)}}{\Sigma_{\beta}^{(1)}} - \sum_{\gamma \in \mathcal{H}} \frac{p_{\alpha\gamma}^{(2)} p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)}}, \quad (\text{B.5})$$

$$M_{\alpha\beta\gamma}^{(3)} = \frac{\partial M_{\alpha\beta}(\boldsymbol{\nu}^{\text{sp}})}{\partial J_{\gamma}} \Big|_{\mathbf{J}=\mathbf{0}} = \sum_{\delta \in \mathcal{H}} F_{\alpha\beta\delta}^{(3)} \Sigma_{\delta\gamma}^{(2)}, \quad (\text{B.6})$$

$$\begin{aligned} F_{\alpha\beta\gamma}^{(3)} &= \frac{p_{\alpha\beta\gamma}^{(3)}}{\Sigma_{\alpha}^{(1)}} + \frac{p_{\alpha\beta\gamma}^{(3)}}{\Sigma_{\beta}^{(1)}} + \frac{p_{\alpha\beta\gamma}^{(3)}}{\Sigma_{\gamma}^{(1)}} - \frac{p_{\alpha\beta}^{(2)} p_{\alpha\gamma}^{(2)}}{\Sigma_{\alpha}^{(1)^2}} - \frac{p_{\alpha\beta}^{(2)} p_{\beta\gamma}^{(2)}}{\Sigma_{\beta}^{(1)^2}} - \frac{p_{\alpha\gamma}^{(2)} p_{\gamma\beta}^{(2)}}{\Sigma_{\gamma}^{(1)^2}} \\ &\quad - \sum_{\delta \in \mathcal{H}} \left(\frac{p_{\alpha\beta\delta}^{(3)} p_{\delta\gamma}^{(2)}}{\Sigma_{\delta}^{(1)}} + \frac{p_{\alpha\delta\gamma}^{(3)} p_{\delta\beta}^{(2)}}{\Sigma_{\delta}^{(1)}} + \frac{p_{\delta\beta\gamma}^{(3)} p_{\alpha\delta}^{(2)}}{\Sigma_{\delta}^{(1)}} - 2 \frac{p_{\alpha\delta}^{(2)} p_{\delta\beta}^{(2)} p_{\delta\gamma}^{(2)}}{\Sigma_{\delta}^{(1)^2}} \right), \end{aligned} \quad (\text{B.7})$$

$$\begin{aligned} F_{\alpha\beta\gamma\delta}^{(4)} &= \frac{p_{\alpha\beta\gamma}^{(3)} p_{\alpha\delta}^{(2)} + p_{\alpha\beta\delta}^{(3)} p_{\alpha\gamma}^{(2)} + p_{\alpha\gamma\delta}^{(3)} p_{\alpha\beta}^{(2)}}{\Sigma_{\alpha}^{(1)^2}} + \frac{p_{\alpha\beta\gamma}^{(3)} p_{\beta\delta}^{(2)} + p_{\alpha\beta\delta}^{(3)} p_{\beta\gamma}^{(2)} + p_{\beta\gamma\delta}^{(3)} p_{\alpha\beta}^{(2)}}{\Sigma_{\beta}^{(1)^2}} \\ &\quad + \frac{p_{\alpha\beta\gamma}^{(3)} p_{\gamma\delta}^{(2)} + p_{\alpha\gamma\delta}^{(3)} p_{\gamma\beta}^{(2)} + p_{\beta\gamma\delta}^{(3)} p_{\alpha\gamma}^{(2)}}{\Sigma_{\gamma}^{(1)^2}} + \frac{p_{\alpha\beta\delta}^{(3)} p_{\delta\gamma}^{(2)} + p_{\alpha\delta\gamma}^{(3)} p_{\delta\beta}^{(2)} + p_{\delta\beta\gamma}^{(3)} p_{\alpha\delta}^{(2)}}{\Sigma_{\delta}^{(1)^2}} \\ &\quad - 2 \left(\frac{p_{\alpha\beta}^{(2)} p_{\alpha\gamma}^{(2)} p_{\alpha\delta}^{(2)}}{\Sigma_{\alpha}^{(1)^3}} + \frac{p_{\alpha\beta}^{(2)} p_{\beta\gamma}^{(2)} p_{\beta\delta}^{(2)}}{\Sigma_{\beta}^{(1)^3}} + \frac{p_{\alpha\gamma}^{(2)} p_{\gamma\beta}^{(2)} p_{\gamma\delta}^{(2)}}{\Sigma_{\gamma}^{(1)^3}} + \frac{p_{\alpha\delta}^{(2)} p_{\delta\beta}^{(2)} p_{\delta\gamma}^{(2)}}{\Sigma_{\delta}^{(1)^3}} \right) \end{aligned}$$

$$\begin{aligned}
& + \sum_{\varepsilon \in \mathcal{H}} \left[\frac{p_{\alpha\beta\varepsilon}^{(3)} p_{\varepsilon\gamma\delta}^{(3)} + p_{\alpha\varepsilon\gamma}^{(3)} p_{\varepsilon\beta\delta}^{(3)} + p_{\varepsilon\beta\gamma}^{(3)} p_{\varepsilon\alpha\delta}^{(3)}}{\Sigma_{\varepsilon}^{(1)}} \right. \\
& \quad - 2 \left(\frac{p_{\alpha\beta\varepsilon}^{(3)} p_{\varepsilon\gamma}^{(2)} p_{\varepsilon\delta}^{(2)} + p_{\alpha\varepsilon\gamma}^{(3)} p_{\varepsilon\beta}^{(2)} p_{\varepsilon\delta}^{(2)} + p_{\varepsilon\beta\gamma}^{(3)} p_{\alpha\varepsilon}^{(2)} p_{\varepsilon\delta}^{(2)}}{\Sigma_{\varepsilon}^{(1)^2}} \right. \\
& \quad \quad \left. + \frac{p_{\alpha\varepsilon\delta}^{(3)} p_{\varepsilon\beta}^{(2)} p_{\varepsilon\gamma}^{(2)} + p_{\varepsilon\beta\delta}^{(3)} p_{\varepsilon\alpha}^{(2)} p_{\varepsilon\gamma}^{(2)} + p_{\varepsilon\gamma\delta}^{(3)} p_{\alpha\varepsilon}^{(2)} p_{\varepsilon\beta}^{(2)}}{\Sigma_{\varepsilon}^{(1)^2}} \right) \\
& \quad \left. + 6 \frac{p_{\varepsilon\alpha}^{(2)} p_{\varepsilon\beta}^{(2)} p_{\varepsilon\gamma}^{(2)} p_{\varepsilon\delta}^{(2)}}{\Sigma_{\varepsilon}^{(1)^3}} \right]. \tag{B.8}
\end{aligned}$$

To find $\mathbf{p}^{(4)}$, we first determine the reduced tensor $\hat{\mathbf{p}}^{(4)}$ which is the solution of the linear system

$$\begin{aligned}
& \sum_{\alpha \in \hat{\mathcal{H}}} \sum_{\beta \in \hat{\mathcal{H}}} \sum_{\gamma \in \hat{\mathcal{H}}} \sum_{\delta \in \hat{\mathcal{H}}} \left(\tilde{\Lambda}_{\alpha_1\alpha}^{(2,0)} \tilde{\Sigma}_{\beta\alpha_2}^{(2)} \tilde{\Sigma}_{\gamma\alpha_3}^{(2)} \tilde{\Sigma}_{\delta\alpha_4}^{(2)} + \tilde{\Sigma}_{\alpha_1\alpha}^{(2,0)} \tilde{\Lambda}_{\beta\alpha_2}^{(2)} \tilde{\Sigma}_{\gamma\alpha_3}^{(2)} \tilde{\Sigma}_{\delta\alpha_4}^{(2)} \right. \\
& \quad \left. + \tilde{\Sigma}_{\alpha_1\alpha}^{(2,0)} \tilde{\Sigma}_{\beta\alpha_2}^{(2)} \tilde{\Lambda}_{\gamma\alpha_3}^{(2)} \tilde{\Sigma}_{\delta\alpha_4}^{(2)} + \tilde{\Sigma}_{\alpha_1\alpha}^{(2,0)} \tilde{\Sigma}_{\beta\alpha_2}^{(2)} \tilde{\Sigma}_{\gamma\alpha_3}^{(2)} \tilde{\Lambda}_{\delta\alpha_4}^{(2)} \right) \hat{p}_{\alpha\beta\gamma\delta}^{(4)} = \Delta_{\alpha_1\alpha_2\alpha_3\alpha_4}, \tag{B.9}
\end{aligned}$$

with $\alpha_1, \alpha_2, \alpha_3, \alpha_4 \in \hat{\mathcal{H}}$. The matrices $\tilde{\Sigma}^{(2)}$, $\tilde{\Sigma}^{(2,0)}$, $\tilde{\Lambda}^{(2)}$ and $\tilde{\Lambda}^{(2,0)}$ are defined by (94), (96), (108) and (109), respectively. The complete fourth order perturbative parameter $\mathbf{p}^{(4)}$ is then recovered using the sum rules (88) for $k = 4$.

References

- [1] De Angelis G F, Jona-Lasinio G and Sirugue M, 1983 *J. Phys. A* **16**, 2433
- [2] De Angelis G F, Jona-Lasinio G and Sidoravicius V, 1998 *J. Phys. A* **31**, 289
- [3] Beccaria M, Presilla C, De Angelis G F and Jona-Lasinio G, 1999 *Europhys. Lett.* **48**, 243
- [4] Beccaria M, Presilla C, De Angelis G F and Jona-Lasinio G, 2000 *Nucl. Phys. B - Proceed. Suppl.* **83-84**, 911-913
- [5] Beccaria M, Presilla C, De Angelis G F and Jona-Lasinio G, 2001 *Int. J. Mod. Phys. B* **15** 1740
- [6] Sorella S, Capriotti L, 2000 *Phys. Rev. B* **61** 2599
- [7] Ostilli M and Presilla C, 2004 *J. Phys. A* **38**, 405
- [8] Ceperley D M, Kalos M H, 1992 *Monte Carlo Methods in Statistical Physics* (Heidelberg: Springer-Verlag).
- [9] Ostilli M and Presilla C, 2004 *New J. Phys.* **6**, 107
- [10] Presilla C and Ostilli M, 2006 *Int. J. Mod. Phys. B* **20**, 2770
- [11] Ostilli M and Presilla C, 2005 *J. Stat. Mech.* P04007
- [12] Ostilli M and Presilla C, 2006 *J. Stat. Mech.* P11012
- [13] Kingman J F C, 1993 *Poisson Processes* (Clarendon Press)
- [14] For $t = 0$ we have $E(\delta_{\mathbf{n}, \mathbf{n}_0} \mathcal{M}_{\mathbf{n}_0}^0) = \langle \mathbf{n}, e^{-i\hat{H}0} \mathbf{n}_0 \rangle$ and for $t > 0$ we calculate $dE(\delta_{\mathbf{n}, \mathbf{n}_{N_t}} \mathcal{M}_{\mathbf{n}_0}^t) = E(\delta_{\mathbf{n}, \mathbf{n}_{N_t+dt}} \mathcal{M}_{\mathbf{n}_0}^{t+dt}) - E(\delta_{\mathbf{n}, \mathbf{n}_{N_t}} \mathcal{M}_{\mathbf{n}_0}^t) = -i \sum_{\mathbf{n}'} \langle \mathbf{n}, \hat{H} \mathbf{n}' \rangle E(\delta_{\mathbf{n}', \mathbf{n}_{N_t}} \mathcal{M}_{\mathbf{n}_0}^t) dt + \mathcal{O}(dt^2)$. Claim follows from the uniqueness of the solution of the system of ordinary differential equations $d\langle \mathbf{n}, e^{-i\hat{H}t} \mathbf{n}_0 \rangle / dt = -i \sum_{\mathbf{n}'} \langle \mathbf{n}, \hat{H} \mathbf{n}' \rangle \langle \mathbf{n}', e^{-i\hat{H}t} \mathbf{n}_0 \rangle$.
- [15] Bremaud P, 1999 *Markov Chains, Gibbs Field, Monte Carlo Simulation, and Queues* (New York: Springer)
- [16] Shiriyayev A N, 1984 *Probability* (New York: Springer)
- [17] Laub A J, 1979 *IEEE Trans. Automat. Control* **24**(6) 913

- [18] Bini D A, Iannazzo B, Meini B, Poloni F, in *Matrix methods: theory, algorithms and applications*, edited by Olshevsky V and Tyrtyshnikov E, World Scientific Publishers, pp 176–209
- [19] Press W H, Teukolsky S A, Vetterling W T, Flannery B P, 1992 *Numerical Recipes, The art of Scientific Computing* 2nd Edition (Cambridge: Cambridge University Press)
- [20] Hubbard J, 1963 *Proc. Roy. Soc. A* **276**, 238
- [21] Giuliani A, Mastropietro V, 2010 *Comm. Math. Phys.* **293**, 301
- [22] Guo C H, 2001 *SIAM J. Matrix Anal. Appl.* **23(1)** 225
- [23] Guo C H, 2006 *J. Comput. Appl. Math.*, **192(2)** 353
- [24] See theorem 4 of [23]
- [25] Anderson E, Bai Z, Bischof C, Blackford S, Demmel J, Dongarra J, Du Croz J, Greenbaum A, Hammarling S, McKenney A, Sorensen D, 1999 *LAPACK Users' Guide* 3rd Edition (Philadelphia: Society for Industrial and Applied Mathematics)